

PCTWORLD INTELLECTUAL PROPERTY ORGANIZATION
International Bureau

INTERNATIONAL APPLICATION PUBLISHED UNDER THE PATENT COOPERATION TREATY (PCT)

(51) International Patent Classification ⁶ : A61K 31/41, 31/36, 31/135		A1	(11) International Publication Number: WO 99/01127
			(43) International Publication Date: 14 January 1999 (14.01.99)
(21) International Application Number: PCT/US98/13807		(81) Designated States: AL, AU, BA, BB, BG, BR, CA, CN, CZ, EE, GE, HU, ID, IL, IS, JP, KP, KR, LC, LK, LR, LT, LV, MG, MK, MN, MX, NO, NZ, PL, RO, SG, SI, SK, SL, TR, TT, UA, US, UZ, VN, YU, ARIPO patent (GH, GM, KE, LS, MW, SD, SZ, UG, ZW), Eurasian patent (AM, AZ, BY, KG, KZ, MD, RU, TJ, TM), European patent (AT, BE, CH, CY, DE, DK, ES, FI, FR, GB, GR, IE, IT, LU, MC, NL, PT, SE), OAPI patent (BF, BJ, CF, CG, CI, CM, GA, GN, ML, MR, NE, SN, TD, TG).	
(22) International Filing Date: 1 July 1998 (01.07.98)			
(30) Priority Data: 60/051,632 3 July 1997 (03.07.97) US			
(71) Applicant (for all designated States except US): SMITHKLINE BEECHAM CORPORATION [US/US]; One Franklin Plaza, Philadelphia, PA 19103 (US).			
(72) Inventors; and (75) Inventors/Applicants (for US only): BONDINELL, William, E. [US/US]; 1512 Franklin Lane, Wayne, PA 19087 (US). CHAN, James, A. [US/US]; 1215 Upton Circle, West Chester, PA 19380 (US).			
(74) Agents: STEIN-FERNANDEZ, Nora et al.; SmithKline Beecham Corporation, Corporate Intellectual Property, UW2220, 709 Swedeland Road, P.O. Box 1539, King of Prussia, PA 19406-0939 (US).		Published <i>With international search report.</i>	
(54) Title: COMPOUNDS AND METHODS			
(57) Abstract <p>This invention relates to substituted benzanilides which are ligands, agonists or antagonists, of the CCR5 receptor. In addition, this invention relates to the treatment and prevention of disease states mediated by CCR5, including, but not limited to, asthma and atopic disorders (for example, atopic dermatitis and allergies), rheumatoid arthritis, sarcoidosis and other fibrotic diseases, atherosclerosis, psoriasis, autoimmune diseases such as multiple sclerosis, and inflammatory bowel disease, all in mammals, by the use of substituted benzanilides which are CCR5 receptor antagonists. Furthermore, since CD8+T cells have been implicated in COPD, CCR5 may play a role in their recruitment and therefore antagonists to CCR5 could provide potential therapeutic in the treatment of COPD. Also, since CCR5 is a co-receptor for the entry of HIV into cells, selective receptor ligands may be useful in the treatment of HIV infection.</p>			

FOR THE PURPOSES OF INFORMATION ONLY

Codes used to identify States party to the PCT on the front pages of pamphlets publishing international applications under the PCT.

AL	Albania	ES	Spain	LS	Lesotho	SI	Slovenia
AM	Armenia	FI	Finland	LT	Lithuania	SK	Slovakia
AT	Austria	FR	France	LU	Luxembourg	SN	Senegal
AU	Australia	GA	Gabon	LV	Latvia	SZ	Swaziland
AZ	Azerbaijan	GB	United Kingdom	MC	Monaco	TD	Chad
BA	Bosnia and Herzegovina	GE	Georgia	MD	Republic of Moldova	TG	Togo
BB	Barbados	GH	Ghana	MG	Madagascar	TJ	Tajikistan
BE	Belgium	GN	Guinea	MK	The former Yugoslav Republic of Macedonia	TM	Turkmenistan
BF	Burkina Faso	GR	Greece	ML	Mali	TR	Turkey
BG	Bulgaria	HU	Hungary	MN	Mongolia	TT	Trinidad and Tobago
BJ	Benin	IE	Ireland	MR	Mauritania	UA	Ukraine
BR	Brazil	IL	Israel	MW	Malawi	UG	Uganda
BY	Belarus	IS	Iceland	MX	Mexico	US	United States of America
CA	Canada	IT	Italy	NE	Niger	UZ	Uzbekistan
CF	Central African Republic	JP	Japan	NL	Netherlands	VN	Viet Nam
CG	Congo	KE	Kenya	NO	Norway	YU	Yugoslavia
CH	Switzerland	KG	Kyrgyzstan	NZ	New Zealand	ZW	Zimbabwe
CI	Côte d'Ivoire	KP	Democratic People's Republic of Korea	PL	Poland		
CM	Cameroon	KR	Republic of Korea	PT	Portugal		
CN	China	KZ	Kazakhstan	RO	Romania		
CU	Cuba	LC	Saint Lucia	RU	Russian Federation		
CZ	Czech Republic	LI	Liechtenstein	SD	Sudan		
DE	Germany	LK	Sri Lanka	SE	Sweden		
DK	Denmark	LR	Liberia	SG	Singapore		
EE	Estonia						

COMPOUNDS AND METHODS

FIELD OF THE INVENTION

5 This invention relates to substituted benzanilides which are ligands, agonists or antagonists, of the CC chemokine receptor CC-CKR5 now designated as CCR5 (*Nature Medicine* **1996**, 2, 1174-8). In addition, this invention relates to the treatment and prevention of disease states mediated by CCR5.

BACKGROUND OF THE INVENTION

10 T cells are not only key regulators of the immune response to infectious agents but are believed critical for the initiation and maintenance of the inflammatory reaction in a variety of chronic diseases. Increased numbers or enhanced activation state of T cells, especially CD4+ T cells, have been demonstrated in the synovium of individuals with rheumatoid arthritis (M.J. Elliott and R. N. Maini, *Int. Arch. Allergy Immunol.* 104: 112-1125, 1994), in the
15 bronchial mucosa of asthmatics (C.J. Corrigan and A.B. Kay, *Immunol. Today* 13:501-506, 1992), in the lesions of multiple sclerosis (R. Martin and H. F. McFarland, *Crit. Rev. Clin. Lab. Sci.* 32: 121-182, 1995), in psoriatic lesions (J.L. Jones, J. Berth-Jone, A. Fletcher and P.E. Hutchinson, *J. Pathol.* 174: 77-82, 1994)
20 and in the fatty streaks of atherosclerosis (R. Ross, *Annu. Rev. Physiol.* 57: 791-804, 1995).

T cells, as well as other inflammatory cells, will migrate into tissues in response to the production of a variety chemotactic factors. Among these factors are a superfamily of 8-12 kDa proteins known as the chemokines. These proteins
25 share structural features such as the presence of 3-4 conserved cysteine residues. RANTES, which stands for Regulated upon Activation Normal T cell Expressed and Secreted, is a 8 kDa protein member of CC branch of the chemokine family. These proteins recruit and activate immune and inflammatory cells through an interaction with G-protein coupled receptors. The CC branch is defined by the
30 absence of an intervening amino acid residue between the first two cysteine residues and members of this family predominately elicit the migration of mononuclear cells, eosinophils and basophils (M. Baggiolini, B. Dewald, and B. Moser, *Adv. Immunol.* 55: 97-179, 1994; and J.J. Oppenheim, C.O.C. Zachariae, N. Mukaida, and K. Matsushima, *Annu. Rev. Immunol.* 9: 617-648, 1991).
35 RANTES potently produces chemotaxis of T cells, basophils, eosinophils, monocytes and mast cells. RANTES was originally identified as gene product induced late after antigen activation of T-cells (T.J. Schall, J. Jongstra, B.J. Dyer, J.

Jorgensen, et al., J. Immunol. 141:1018-1025, 1988), however, RANTES has been shown to be synthesized and secreted by a diverse group of cells that include epithelial and endothelial cells (C. Stellato, L.A. Beck, G.A. Gorgone, D. Proud, et al., J. Immunol. 155: 410-418, 1995; and A. Marfaing-Koka, O. Devergne, G. Gorgone, A. Portier, et al., J. Immunol. 154: 1870-1878, 1994), synovial fibroblasts (P. Rathanaswami, M. Hachicha, M. Sadick, T.J. Schall, et al., J. Biol. Chem. 268: 5834-5839, 1993) and dermal fibroblasts (M. Sticherling, M. Kupper, F. Koltowitz, E. Bornscheuer, et al., J. Invest. Dermatol. 105: 585-591, 1995), mesangial cells (G. Wolf, S. Aberle, F. Thaiss, et al., Kidney Int. 44: 795-804, 1994) and platelets (Y. Koameyoshi, A. Dorschner, A.I. Mallet, E. Christophers, et al., J. Exp. Med. 176: 587-592, 1992). In these cells RANTES mRNA is rapidly upregulated in response to IL-1 or TNF α . Although RANTES mRNA is not usually detected in normal tissues (J.M. Pattison, P.J. Nelson, and A.M. Krensky, Clin. Immunother. 4: 1-8, 1995), increased mRNA or protein has been found in diseases characterized by a mononuclear infiltrate. For example, RANTES mRNA was visualized using *in situ* hybridization in renal allografts undergoing rejection (J.M. Pattison, P.J. Nelson, and A.M. Krensky, Clin. Immunother. 4: 1-8, 1995; and K.C. Nadeau, H. Azuma and N.I. Tilney, Proc. Natl. Acad. USA 92: 8729-8733, 1995) in the skin of atopic dermatitis patients after exposure to antigen (S. Ying, L. Taborda-Barata, Q. Meng, M. Humbert, et al., J. Exp. Med. 181: 2153-2159, 1995), and in endothelial cells of coronary arteries undergoing accelerated atherosclerosis after cardiac transplant (J.M. Pattison, P.J. Nelson, and A.M. Krensky, Clin. Immunother. 4: 1-8, 1995). Further, increased immunoreactive protein for RANTES has been detected in bronchoalveolar lavage fluid (R. Alam, J. York, M. Boyers, et al., Am. J. Resp. Crit. Care Med. 149: A951, 1994) and sputum from asthmatic individuals (C.M. Gelder, P.S. Thomas, D.H. Yates, I.M. Adcock, et al., Thorax 50: 1033-1037, 1995).

Several receptors have been identified that bind RANTES. In particular, CCR5, when expressed in either HEK 293 cells or CHO cells, binds RANTES. This receptor is expressed in T-cells and in monocytes and macrophages, immune/inflammatory cells which are important in the maintenance of a chronic inflammatory reaction. Pharmacological characterization of CCR5 indicates similarities to the RANTES binding site observed on isolated T cells. Therefore, antagonism of RANTES' action on CCR5, as well as antagonism of other natural ligands of CCR5, should inhibit the recruitment of T cells into inflammatory lesions and provide a novel therapeutic approach for the treatment of atopic and autoimmune disorders.

Since T cells express CCR5, selective receptor ligands of CCR5, particularly antagonists, are likely to provide beneficial effects in diseases including, but not limited to, asthma and atopic disorders (for example, atopic dermatitis and allergies), rheumatoid arthritis, sarcoidosis and other fibrotic diseases, atherosclerosis, psoriasis, autoimmune diseases such as multiple sclerosis, and inflammatory bowel disease, all in mammals, preferably humans. Furthermore, since CD8+ T cells have been implicated in COPD, CCR5 may play a role in their recruitment and therefore antagonists to CCR5 could provide potential therapeutic in the treatment of COPD. Also, since CCR5 is a co-receptor for the entry of HIV into cells, selective receptor ligands may be useful in the treatment of HIV infection.

A subset of compounds included in formula (I) have been reported to have 5-HT receptor activity (international application publication number WO 95/15954, published 15 June 1995, international application publication number WO 95/17398, published 29 June 1995, international application publication number WO 95/26328, published 5 October 1995, international application publication number WO 96/06079, published 29 February 1996, GB 2276161 published 21 September 1994, and GB 2276165 published 21 September 1994; international application publication number WO 95/30675 published 16 November 1995; international application publication number WO 95/17401 published 29 June 1995; international application publication number WO 96/31508 published 10 October 1996; international application publication number WO 97/10824 published 27 March 1997; international application publication number WO 96/11934 published 25 April 1996; international application publication number WO 96/19477 published 27 June 1996; international application publication number WO 97/17350 published 15 May 1997; international application publication number WO 97/34900 published 25 September 1997; international application publication number WO 97/34901 published 25 September 1997; international application publication number WO 97/35862 published 2 October 1997; international application publication number WO 97/19070 published 29 May 1997; international application publication number WO 95/32967 published 7 December 1995; international application publication number WO 97/07120 published 27 February 1997; U.S. Patent 3,931,195, issued Jan. 6, 1976; and U.S. Patent 4,000,143, issued Dec. 28, 1976).

Surprisingly, it has now been discovered that this class of non-peptide compounds, in particular substituted benzanilides of formula (I), function as CCR5

receptor ligands, and therefore, have utility in the treatment and prevention of disease states mediated by CCR5 receptor mechanisms.

SUMMARY OF THE INVENTION

5 The present invention is to the novel use of a CCR5 ligand for the treatment of certain disease states, including, but not limited to, asthma and atopic disorders (for example, atopic dermatitis and allergies), rheumatoid arthritis, sarcoidosis and other fibrotic diseases, atherosclerosis, psoriasis, autoimmune diseases such as multiple sclerosis, inflammatory bowel disease, and HIV infection, all in
10 mammals, preferably humans. The preferred compounds for use as CCR5 ligands are those compounds of Formula (I) as noted herein.

DETAILED DESCRIPTION OF THE INVENTION

 It has now been discovered that substituted of formula (I) are CCR5
15 receptor ligands. It has also now been discovered that selective inhibition of CCR5 receptor mechanisms by treatment with the receptor ligands of formula (I), or a pharmaceutically acceptable salt thereof, represents a novel therapeutic and preventative approach to the treatment of a variety of disease states, including, but not limited to, asthma and atopic disorders (for example, atopic dermatitis and
20 allergies), rheumatoid arthritis, sarcoidosis and other fibrotic diseases, atherosclerosis, psoriasis, autoimmune diseases such as multiple sclerosis, and inflammatory bowel disease, all in mammals, preferably humans. Furthermore, since CD8+ T cells have been implicated in COPD, CCR5 may play a role in their recruitment and therefore antagonists to CCR5 could provide potential therapeutic
25 in the treatment of COPD. Also, since CCR5 is a co-receptor for entry into cells, selective receptor ligands may be useful in the treatment of HIV infection.

 Compounds for use herein include the CCR5 ligands as described in international application publication number WO 95/15954, published 15 June 1995, (USSN 08/652,581, filed June 7, 1996); international application publication
30 number WO 95/17398, published 29 June 1995, (USSN 08/663,291, filed June 21, 1996); international application publication number WO 95/26328, published 5 October 1995, (USSN 08/718,481, filed Sept. 26, 1996); international application publication number WO 96/06079, published 29 February 1996, (USSN 08/793,428, filed Feb. 21, 1997); GB 2276161 published 21 September 1994, and
35 GB 2276165 published 21 September 1994; international application publication number WO 95/30675, published 16 November 1995, (USSN 08/737,147); international application publication number WO 95/17401, published 29 June


1995 (USSN 08/663,290, filed June 21, 1996); international application publication number WO 96/31508, published 10 October 1996 (USSN 08/930,848, filed Oct. 7, 1997); international application publication number WO 97/10824, published 27 March 1997 (USSN 09/043,346); international application
 5 publication number WO 96/11934, published 25 April 1996, (USSN 08/817,619); international application publication number WO 96/19477, published 27 June 1996 (USSN 08/849,932); international application publication number WO 97/17350, published 15 May 1997 (USSN 09/068,382, filed May 8, 1998); international application publication number WO 97/34900, published 25
 10 September 1997 (Attorney Docket No. P31419); international application publication number WO 97/34901, published 25 September 1997 (Attorney Docket No. P31418); international application publication number WO 97/35862, published 2 October 1997 (Attorney Docket No. P31426); international application publication number WO 97/19070, published 29 May 1997 (USSN 09/077,263);
 15 international application publication number WO 95/32967, published 7 December 1995 (USSN 08/737,660); international application publication number WO 97/07120, published 27 February 1997 (USSN 09/011,338, filed Feb. 11, 1998); U.S. Patent 3,931,195, issued Jan. 6, 1976; and U.S. Patent 4,000,143, issued Dec. 28, 1976.

20 Preferred compounds for use as CCR5 ligands are those compounds of Formula (I) as noted herein.

Each of these references is incorporated herein in their entirety.


A preferred group of compounds for use herein are those compounds of the formula (I) or a pharmaceutically acceptable salt thereof:

25



Formula I

in which Ar represents a group selected from (i), (ii) or (iii);



in which:

30

R^1 and R^2 are independently hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl,

C_{2-6} alkynyl, C_{3-7} cycloalkyl, C_{3-6} cycloalkenyl, aryl, $(CH_2)_aNR^7R^8$,
 $(CH_2)_aNR^7COR^9$, $(CH_2)_aNR^7CO_2R^{10}$, $(CH_2)_aNR^7SO_2R^{11}$,
 $(CH_2)_aCONR^{12}R^{13}$, hydroxy C_{1-6} alkyl, C_{1-4} alkoxyalkyl (optionally substituted
 by a C_{1-4} alkoxy or hydroxy group), $(CH_2)_aCO_2C_{1-6}$ alkyl, $(CH_2)_bOC(O)R^{14}$,
 5 $CR^{15}=NOR^{16}$, $CNR^{15}=NOR^{16}$, COR^{17} , $CONR^{12}R^{13}$, $CONR^{12}(CH_2)_cOC_{1-4}$
 alkyl, $CONR^{12}(CH_2)_aCO_2R^{18}$, $CONHNR^{19}R^{20}$, $CONR^{12}SO_2R^{21}$, CO_2R^{22} ,
 cyano, trifluoromethyl, NR^7R^8 , NR^7COR^9 , $NR^{23}CO(CH_2)_aNR^{23}R^{24}$,
 $NR^{23}CONR^{23}R^{24}$, $NR^7CO_2R^{10}$, $NR^7SO_2R^{11}$, $N=CNR^{23}NR^{23}R^{24}$, nitro,
 hydroxy, C_{1-6} alkoxy, hydroxy C_{1-6} alkoxy, C_{1-6} alkoxy C_{1-6} alkoxy,
 10 $OC(O)NR^{25}R^{26}$, SR^{27} , SOR^{28} , SO_2R^{28} , $SO_2NR^{25}R^{26}$ or halogen;

a is 1, 2, 3 or 4;

R^7 and R^8 are independently hydrogen or C_{1-6} alkyl, or NR^7R^8 forms a
 heterocyclic ring which has 5 or 6 ring members which, may optionally be
 substituted by an oxo group and, when there are 6 ring members, may optionally
 15 contain in the ring one oxygen or sulfur atom;

R^9 is hydrogen, C_{1-6} alkyl or C_{1-4} alkoxyalkyl;

R^{10} is C_{1-6} alkyl;

R^{11} is C_{1-6} alkyl or phenyl;

R^{12} and R^{13} are independently hydrogen or C_{1-6} alkyl, or $NR^{12}R^{13}$ forms
 20 a saturated heterocyclic ring which has 5 or 6 members which, when there are 6
 ring members, may optionally contain in the ring one oxygen or sulfur atom;

b is 0, 1, 2 or 3;

R^{14} is C_{1-4} alkyl, optionally substituted by a C_{1-6} alkoxy;

R^{15} and R^{16} are independently hydrogen or C_{1-6} alkyl;

25 R^{17} is hydrogen or C_{1-6} alkyl;

c is 1, 2 or 3;

R^{18} is hydrogen or C_{1-6} alkyl;

R^{19} and R^{20} are independently hydrogen or C_{1-6} alkyl;

R^{21} is hydrogen or C_{1-6} alkyl;

30 R^{22} is hydrogen or C_{1-6} alkyl optionally substituted with one or two
 substituents selected from C_{1-6} alkyl, C_{1-6} alkoxy, hydroxy, or NR^7R^8 ;

R^{23} and R^{24} are independently hydrogen or C_{1-6} alkyl;

R^{25} and R^{26} are independently hydrogen or C_{1-6} alkyl, or $NR^{25}R^{26}$ forms
 a saturated heterocyclic ring which has 5 or 6 members which, when there are 6
 35 ring members, may optionally contain in the ring one oxygen or sulfur atom;

R^{27} is hydrogen or C_{1-6} alkyl;

R^{28} is C_{1-6} alkyl;

P is a 5 to 7-membered heterocyclic ring containing 1 to 4 heteroatoms selected from oxygen, nitrogen or sulfur;

R^3 and R^4 are independently hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl, C_{3-6} cycloalkenyl, hydroxy C_{1-6} alkyl, C_{1-6} alkylOC $_{1-6}$ alkyl, CONR²⁹R³⁰, CO₂R³¹, cyano, aryl, trifluoromethyl, NR²⁹R³⁰, nitro, hydroxy, C_{1-6} alkoxy, acyloxy or halogen;

R^{29} , R^{30} and R^{31} are independently hydrogen or C_{1-6} alkyl;

R^5 is hydrogen, C_{1-6} alkyl, C_{1-6} alkoxy or halogen;

R^6 is hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl (optionally substituted by a hydroxy or an oxo group), hydroxy C_{1-6} alkyl, hydroxy C_{3-6} alkenyl, hydroxy C_{3-6} alkynyl, $(CH_2)_dOR^{32}$, $(CH_2)_dCOR^{33}$, $(CH_2)_dCR^{34}=NOR^{35}$, CONR³⁶R³⁷, CO₂R³⁸, hydroxy, O $(CH_2)_eR^{39}$, NR³⁶R³⁷, SR⁴⁰, SO₂NR⁴¹R⁴² or halogen; d is 0, 1, 2, 3, 4, 5, or 6;

R^{32} is C_{1-6} alkyl, hydroxy C_{1-6} alkyl, or C_{1-4} alkanoyl;

R^{33} is hydrogen or C_{1-6} alkyl;

R^{34} is hydrogen or C_{1-6} alkyl;

R^{35} is hydrogen or C_{1-6} alkyl;

R^{36} and R^{37} are independently hydrogen or C_{1-6} alkyl or NR³⁶R³⁷ forms a saturated heterocyclic ring which has 5 or 6 members, may optionally be substituted by an oxo group and, when there are 6 ring members, may optionally contain one oxygen or sulfur atom or an NH or NR⁴³ where R^{43} is C_{1-6} alkyl, COR⁴⁴ or CO₂R⁴⁵ where R^{44} and R^{45} are independently hydrogen or C_{1-6} alkyl;

R^{38} is hydrogen or C_{1-6} alkyl;

e is 1, 2, 3, 4, 5 or 6;


R^{39} is C_{1-6} alkoxy, CO₂H, CO₂ C_{1-6} alkyl or CONR³⁶R³⁷;

R^{40} is C_{1-6} alkyl;

R^{41} and R^{42} are independently hydrogen or C_{1-6} alkyl;

alternatively, R^5 and R^6 form a fused benzo ring optionally substituted with C_{1-6} alkyl, C_{1-6} alkoxy or halogen;

when Ar is (i), (ii) or (iii), and A is CONR⁴⁶, NHCO, -NHCH₂, or CH₂NH, where R^{46} is hydrogen or C_{1-6} alkyl, E represents (a):



in which

R^{47} and R^{48} are independently hydrogen or C_{1-6} alkyl;

R^{49} is hydrogen, C_{1-6} alkyl, CO_2R^{53} , $NHCO_2R^{54}$, hydroxy, C_{1-6} alkoxy or halogen where R^{53} is hydrogen or C_{1-6} alkyl and R^{54} is C_{1-6} alkyl;

R^{50} and R^{51} are independently hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl, aralkyl, or together with the nitrogen atom to which they are attached form an optionally substituted 5- to 7-membered heterocyclic ring containing one to two heteroatoms selected from oxygen, nitrogen or sulfur;


B is oxygen, $S(O)_h$ where h is 0, 1 or 2, $CR^{55}=CR^{56}$ or $CR^{55}R^{56}$ where R^{55} and R^{56} are independently hydrogen or C_{1-6} alkyl, or B is NR^{57} where R^{57} is hydrogen, C_{1-6} alkyl or phenyl C_{1-6} alkyl;

R^{52} is hydrogen or R^{52} taken together with R^{46} forms a group D where D is $(CR^{58}R^{59})_i$ where i is 2, 3 or 4 and R^{58} and R^{59} are independently hydrogen or C_{1-6} alkyl or D is $(CR^{58}R^{59})_j-G$ where j is 0, 1, 2 or 3 and G is oxygen, sulfur or $CR^{58}=CR^{59}$;

f is 1 to 4; and

g is 1 or 2;

when Ar is (i), (ii) or (iii), and A is $CONR^{60}$, $NHCO$, or CH_2NH , where R^{60} is hydrogen or C_{1-6} alkyl, E represents (b):



R^{61} is hydrogen or C_{1-6} alkyl or R^{61} and R^{60} together form a group -K- where K is $(CR^{65}R^{66})_m$ where m is 2, 3, or 4 and R^{65} and R^{66} are independently hydrogen or C_{1-6} alkyl or K is $(CR^{65}R^{66})_n-L$ where n is 0, 1, 2, or 3 and L is oxygen, sulfur or $CR^{65}=CR^{66}$;

R^{62} is hydrogen or C_{1-6} alkyl;


R^{63} and R^{64} are independently hydrogen or C_{1-6} alkyl;

J is oxygen, $CR^{67}R^{68}$ or NR^{69} where R^{67} , R^{68} and R^{69} are independently hydrogen or C_{1-6} alkyl or J is a group $S(O)_m$ where m is 0, 1 or 2;

k is 1, 2 or 3;

l is 1, 2 or 3;

when Ar is (i), (ii) or (iii), and A is $CONR^{70}$, $NHCO$, $-NHCH_2$, or CH_2NH , where R^{70} is hydrogen or C_{1-6} alkyl, E represents (c):



in which:

M is oxygen, $S(O)_p$ where p is 0, 1 or 2, $CR^{76}=CR^{77}$ or $CR^{76}R^{77}$ where R^{76} and R^{77} are independently hydrogen or C_{1-6} alkyl, or M is NR^{78} where R^{78} is hydrogen or alkyl;

5 R^{71} and R^{72} are independently hydrogen or C_{1-6} alkyl;

R^{73} is hydrogen, C_{1-6} alkyl, CO_2R^{79} , $NHCO_2R^{80}$, hydroxy, C_{1-6} alkoxy or halogen where R^{79} is hydrogen or C_{1-6} alkyl and R^{80} is C_{1-6} alkyl;


R^{74} is hydrogen or together with R^{70} forms a group Q where Q is $CR^{81}=CR^{82}$, $CR^{81}=CR^{82}CR^{81}R^{82}$ or $(CR^{81}R^{82})_q$ where q is 2 or 3 and R^{81}

10 and R^{82} are independently hydrogen or C_{1-6} alkyl;

n is 0, 1, 2 or 3;


o is 1 or 2;

R^{75} is a group of formula (d):



15 where r, s and t are independently integers having the value 1, 2 or 3;


or R^{75} is a group of formula (e):



where u is 0, 1, 2 or 3 and R^{83} is hydrogen or C_{1-6} alkyl;

when Ar is (i), (ii), or (iii), and A is CONH, NHCO or CH_2NH , E

20 represents (f):




R^{84} and R^{85} are independently hydrogen or C_{1-6} alkyl;

R^{86} and R^{87} are independently hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl, aralkyl, or together with the nitrogen atom to which they are attached form an optionally substituted 5- to 7-membered heterocyclic ring containing one to two heteroatoms selected from oxygen, nitrogen or sulfur;

T is $-(CR^{88}R^{89})_w-$ or $-O(CR^{88}R^{89})_x-$ where R^{88} and R^{89} are independently hydrogen or C_{1-6} alkyl where w is 2 or 3 and x is 1, 2 or 3; v is 1 to 4; and

W is oxygen, $S(O)_y$ where y is 0, 1 or 2, or W is NR^{90} where R^{90} is hydrogen or C_{1-6} alkyl, or W is $CR^{91}=CR^{92}$ or $CR^{91}R^{92}$ where R^{91} and R^{92} are independently hydrogen or C_{1-6} alkyl;

when Ar is (i), (ii) or (iii), and A is $CONR^{93}$, $NHCO$, or CH_2NH where R^{93} is hydrogen or C_{1-6} alkyl, E represents a group (g):




R^{94} is hydrogen, halogen, hydroxy, C_{1-6} alkyl or C_{1-6} alkoxy or R^{94} and R^{93} taken together from a group $-X-$ where X is $(CR^{97}R^{98})_{aa}$ where aa is 2, 3 or 4 and R^{97} and R^{98} are independently hydrogen or C_{1-6} alkyl or X is $(CR^{97}R^{98})_{ab}-Y$ where ab is 0, 1, 2 or 3 and Y is oxygen, sulfur or $CR^{97}=CR^{98}$;

R^{95} is hydrogen, C_{1-6} alkyl, CO_2R^{99} , $NHCO_2R^{100}$, hydroxy, C_{1-6} alkoxy or halogen where R^{95} is hydrogen or C_{1-6} alkyl and R^{100} is C_{1-6} alkyl; z is 1 or 2; and

R^{96} is an optionally substituted 5 to 7-membered saturated or partially saturated heterocyclic ring containing 1 to 3 heteroatoms selected from nitrogen, oxygen or sulfur or R^{96} is an optionally substituted 6,6 or 6,5 bicyclic ring containing a nitrogen atom and optionally a further heteroatom selected from oxygen, nitrogen or sulfur;

when Ar is (i), (ii) or (iii) and A is $CONR^{101}$, $NHCO$, or CH_2NH , where R^{101} is hydrogen or C_{1-6} alkyl, E represents group (h):



R^{102} is hydrogen or C_{1-6} alkyl, C_{1-6} alkoxy or halogen, or R^{102} together with R^{101} form a group $-AA-$ where AA is $(CR^{107}R^{108})_{ad}$ where ad is 1, 2 or 3 and R^{107} and R^{108} are independently hydrogen or C_{1-6} alkyl or AA is $(CR^{107}=CR^{108})_{ae}-AB$ where ae is 0, 1 or 2 and AB is oxygen, sulfur, $CR^{107}=CR^{108}$, $CR^{107}=N$, $CR^{102}NR^{108}$ or $N=N$;

R^{103} and R^{104} are independently hydrogen or C_{1-6} alkyl;


R^{105} and R^{106} are independently hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl, aralkyl, or together with the nitrogen atom to which they are attached form an

optionally substituted 5- to 7-membered heterocyclic ring containing one to two heteroatoms selected from oxygen, nitrogen or sulfur;

ac is 0 to 4;

5 Z is an optionally substituted 5 to 7-membered heterocyclic ring containing 1 to 3 heteroatoms selected from oxygen, nitrogen or sulfur;

when Ar is (i), (ii) or (iii) and A is CONR¹⁰⁹, NHCO, or CH₂NH where R¹⁰⁹ is hydrogen or C₁₋₆alkyl, E represents group (i):



(i);

R¹¹⁰ is hydrogen or C₁₋₆alkyl or R¹¹⁰ and R¹⁰⁹ together form a group - AD- where AD is (CR¹¹⁵R¹¹⁶)_{ah} where ah is 2, 3 or 4 and R¹¹⁵ and R¹¹⁶ are independently hydrogen or C₁₋₆alkyl or AD is (CR¹¹⁵R¹¹⁶)_{ai}-AE where ai is 0, 1, 2 or 3 and AE is oxygen, sulfur or CR¹¹⁵=CR¹¹⁶;

R¹¹¹ and R¹¹² are independently hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl, aralkyl, or together with the nitrogen atom to which they are attached form an optionally substituted 5- to 7-membered heterocyclic ring containing one to two heteroatoms selected from oxygen, nitrogen or sulfur;

R¹¹³ and R¹¹⁴ are independently hydrogen or C₁₋₆alkyl;

AC is oxygen, CR¹¹⁷R¹¹⁸ or NR¹¹⁹ where R¹¹⁷, R¹¹⁸ and R¹¹⁹ are independently hydrogen or C₁₋₆alkyl or AC is a group S(O)_{aj} where aj is 0, 1 or 2; af is 1, 2 or 3;

ag is 1, 2, 3, or 4; and

ah is 0, 1, 2, 3 or 4.

Suitably, when Ar is (i) or (ii), the terminal phenyl group in (i) and (ii) can be attached to the phenyl group bearing group A in any position. Preferably the terminal phenyl ring is attached to the phenyl bearing group A in a position meta or para to group A, more preferably para to group A.

Suitably P is a 5- to 7-membered heterocyclic ring containing 1 to 4 heteroatoms selected from oxygen, nitrogen or sulfur. Examples of suitable heterocyclic rings include thienyl, furyl, triazolyl, diazolyl, imidazolyl, oxadiazolyl, isothiazolyl, isoxazolyl, thiadiazolyl, pyridyl, pyrimidyl, pyrazinyl and dioxanyl. Suitably the heterocyclic rings can be linked to the remainder of the molecule via a carbon atom or, when present, a nitrogen atom. Preferably P is oxadiazolyl.

Suitably, when Ar is (iii), R⁶ can be attached to the phenyl ring in any position. Preferably, R⁶ is attached meta or para to group A. More preferably, R⁶ is attached at the para to group A.

Suitably R¹ and R² are independently hydrogen, C₁₋₆alkyl, C₂₋₆alkenyl, C₂₋₆alkynyl, C₃₋₇cycloalkyl, C₃₋₆cycloalkenyl, aryl, (CH₂)_aNR⁷R⁸, (CH₂)_aNR⁷COR⁹, (CH₂)_aNR⁷CO₂R¹⁰, (CH₂)_aNR⁷SO₂R¹¹, (CH₂)_aCONR¹²R¹³, hydroxyC₁₋₆alkyl, C₁₋₄alkoxyalkyl (optionally substituted by a C₁₋₄alkoxy or hydroxy group), (CH₂)_aCO₂C₁₋₆alkyl, (CH₂)_bOC(O)R¹⁴, CR¹⁵=NOR¹⁶, CNR¹⁵=NOR¹⁶, COR¹⁷, CONR¹²R¹³, CONR¹²(CH₂)_cOC₁₋₄alkyl, CONR¹²(CH₂)_aCO₂R¹⁸, CONHN¹⁹R²⁰, CONR¹²SO₂R²¹, CO₂R²², cyano, trifluoromethyl, NR⁷R⁸, NR⁷COR⁹, NR²³CO(CH₂)_aNR²³R²⁴, NR²³CONR²³R²⁴, NR⁷CO₂R¹⁰, NR⁷SO₂R¹¹, N=CNR²³NR²³R²⁴, nitro, hydroxy, C₁₋₆alkoxy, hydroxyC₁₋₆alkoxy, C₁₋₆alkoxyC₁₋₆alkoxy, OC(O)NR²⁵R²⁶, SR²⁷, SOR²⁸, SO₂R²⁸, SO₂NR²⁵R²⁶ or halogen. Preferably, at least one of R¹ or R² are independently, hydrogen, C₁₋₆alkyl, preferably methyl, C₁₋₆alkoxy, preferably methoxy, preferably in the 3-position, hydroxy, preferably in the 3-position, cyano, preferably in the 4-position, and (CH₂)_bOC(O)R¹⁴, wherein b is preferably 0, and R¹⁴ is preferably ethyl.

Suitably R³ and R⁴ are independently hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl, C₃₋₆cycloalkenyl, hydroxyC₁₋₆alkyl, C₁₋₆alkylOC₁₋₆alkyl, CONR²⁹R³⁰, CO₂R³¹, cyano, aryl, trifluoromethyl, NR²⁹R³⁰, nitro, hydroxy, C₁₋₆alkoxy, acyloxy or halogen. Preferably R³ is hydrogen and C₁₋₆alkyl, more preferably methyl. R⁴-P is preferably CH₃-oxadiazolyl.

Suitably R⁶ is hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl (optionally substituted by a hydroxy or an oxo group), hydroxyC₁₋₆alkyl, hydroxyC₃₋₆alkenyl, hydroxyC₃₋₆alkynyl, (CH₂)_dOR³², (CH₂)_dCOR³³, (CH₂)_dCR³⁴=NOR³⁵, CONR³⁶R³⁷, CO₂R³⁸, hydroxy, O(CH₂)_eR³⁹, NR³⁶R³⁷, SR⁴⁰, SO₂NR⁴¹R⁴² or halogen. Preferably, R⁶ is C₃₋₇cycloalkyl, more preferably cyclohexyl, or halogen, more preferably iodo.

Preferably R⁷ and R⁸ are independently hydrogen or C₁₋₆alkyl;
 Preferably R⁹ is hydrogen or C₁₋₆alkyl;
 Preferably R¹¹ is C₁₋₆alkyl;
 Preferably R¹² and R¹³ are independently hydrogen or C₁₋₆alkyl;
 Preferably R¹⁴ is C₁₋₆alkyl;
 Preferably R²² is hydrogen or C₁₋₆alkyl;
 Preferably R²⁵ and R²⁶ are independently hydrogen or C₁₋₆alkyl.
 Suitably E represents (a). When E represents (a), the groups

-B(CR⁴⁷R⁴⁸)_f-NR⁵⁰R⁵¹ and R⁴⁹ can be attached to the phenyl ring at any position. Preferably the group -B(CR⁴⁷R⁴⁸)_f-NR⁵⁰R⁵¹ is located meta or para to the amide linkage, more preferably meta. Preferably the group R⁴⁹ is located para to the amide linkage. Preferably the group R⁴⁹ is alkoxy, more preferably methoxy, or halogen, more preferably iodo. Preferably, B is oxygen or CR⁵⁵R⁵⁶.
5 Preferably, f is 2 or 3. Suitably, R⁵⁰ and R⁵¹ are C₁₋₆alkyl, preferably isopropyl, or tert-butyl; C₃₋₇cycloalkyl, preferably cyclohexyl; or together with the nitrogen atom to which they are attached, form a 5- to 7-membered heterocyclic ring.

Preferred compounds of formula (I) are wherein, E represents (a), and
10 wherein B is oxygen, CR⁵⁵R⁵⁶ is CH₂ and f is 2, and wherein B is CH₂, CR⁵⁵R⁵⁶ is CH₂ and f is 2, and wherein B is oxygen, CR⁵⁵R⁵⁶ is CH₂ and f is 3. Particularly preferred is a compound of formula (I) wherein B is oxygen, CR⁵⁵R⁵⁶ is CH₂ and f is 2. R⁵² is preferably hydrogen.

Other preferred compounds of formula (I) are wherein, E represents (a) and
15 preferably R⁵⁰ and R⁵¹ are isopropyl, R⁵⁰ is isopropyl and R⁵¹ is tert-butyl or cyclohexyl, or NR⁵⁰R⁵¹ is 2,2,6,6-tetramethylpiperidine. More preferably, R⁵⁰ and R⁵¹ are isopropyl, R⁵⁰ is isopropyl and R⁵¹ is tert-butyl, or NR⁵⁰R⁵¹ is 2,2,6,6-tetramethylpiperidine. Preferably R⁴⁹ is methoxy.

When E represents (b), k is preferably 1, J is preferably oxygen, and R⁶² is
20 preferably C₁₋₆alkyl, more preferably methyl or isopropyl.

When E represents (c), n is preferably 1, M is preferably oxygen, R⁷¹ and R⁷² are preferably hydrogen, R⁷³ is preferably para-alkoxy, more preferably para-OCH₃, R⁷⁴ is preferably hydrogen, and R⁷⁵ is preferably a group of formula (e) wherein u is 2, and R⁸³ is preferably C₁₋₆alkyl, more preferably methyl.

25 The term "C₁₋₆alkyl" is used herein at all occurrences to mean a straight or branched chain radical of 1 to 6 carbon atoms, unless the chain length is limited thereto, including, but not limited to methyl, ethyl, n-propyl, isopropyl, n-butyl, sec-butyl, isobutyl, tert-butyl, and the like.

The terms "halo" or "halogen" are used interchangeably herein at all
30 occurrences to mean radicals derived from the elements chlorine, fluorine, iodine and bromine.

The terms "cycloalkyl" and "cyclic alkyl" are used herein at all occurrences to mean cyclic radicals, preferably comprising 3 to 7 carbon atoms which may be mono- or bicyclo- fused ring systems which may additionally include unsaturation,
35 including, but not limited to, cyclopropyl, cyclopentyl, cyclohexyl, 1,2,3,4-tetrahydronaphthyl, and the like.

The term "alkenyl" is used herein at all occurrences to mean a straight or branched chain radical of 2 to 6 carbon atoms, unless the length is limited thereto, wherein there is at least one double bond between two of the carbon atoms in the chain, including, but not limited to, ethenyl, 1-propenyl, 2-propenyl, 2-methyl-1-propenyl, 1-butenyl, 2-butenyl, and the like.

The term "cycloalkenyl" is used herein to mean cyclic radicals, preferably of 5 to 8 carbons, which have at least one double bond between two of the carbon atoms in the ring, including but not limited to cyclopentenyl, cyclohexenyl, and the like.

The term "alkynyl" is used herein at all occurrences to mean a straight or branched chain radical of 2 to 8 carbon atoms, unless the chain length is limited thereto, wherein there is at least one triple bond between two of the carbon atoms in the chain, including, but not limited to, acetylene, 1-propylene, 2-propylene, and the like.

The term "aryl" is used herein at all occurrences to mean 5-14-membered substituted or unsubstituted aromatic ring(s) or ring systems which may include bi- or tri-cyclic systems, including, but not limited to phenyl, naphthyl, and the like.

The term "aralkyl" is used herein at all occurrences to mean an aryl moiety as defined above, which is connected to an alkyl moiety as defined above, for example, benzyl or phenethyl, and the like.

The term "alkoxy" is used herein at all occurrences to mean a straight or branched chain radical of 1 to 6 carbon atoms, unless the chain length is limited thereto, bonded to an oxygen atom, including, but not limited to, methoxy, ethoxy, n-propoxy, isopropoxy, and the like.

The terms "hydroxyC₁₋₆alkyl" and "hydroxyalkyl" are used herein interchangeably to mean an hydroxyl group bonded to a C₁₋₆alkyl group as defined above, including, but not limited to methanol, ethanol, n-propanol, isopropanol, n-butanol, sec-butanol, isobutanol, tert-butanol, and the like.

The term "C₁₋₄alkoxyalkyl" is used herein at all occurrences to mean a C₁₋₄alkoxy group as defined above bonded to an alkyl group as defined above, such as an ether, e.g., CH₃-CH₂-O-CH₂-CH₂-CH₃.

The term "hydroxyC₁₋₆alkoxy" is used herein at all occurrences to mean an hydroxyl group bonded to an alkoxy group as defined above, e.g., HO-CH₂-CH(OH)CH₃.

The term "C₁₋₆alkoxyC₁₋₆alkoxy" is used herein at all occurrences to mean an alkoxy group as defined above, substituted with an alkoxy group as defined above.

The term "acyloxy" is used herein at all occurrences to mean a moiety -O-C(O)-R, wherein R is hydrogen or C₁₋₆alkyl.

The term "C₁₋₄alkanoyl" is used herein at all occurrences to mean a C(O)C₁₋₄alkyl group wherein the alkyl portion is as defined above.

5 The term "heteroatom" is used herein at all occurrences to mean an oxygen atom, a sulfur atom or a nitrogen atom. It will be recognized that when the heteroatom is nitrogen, it may form an NR_aR_b moiety, wherein R_a and R_b are, independently, hydrogen or C₁ to C₆ alkyl, or together with the nitrogen to which they are bound, form a saturated or unsaturated 5-, 6- or 7-membered ring, including, but not limited to, pyrrolidine, piperidine, piperazine, morpholine, 10 pyridine, and the like. It will be recognized that the saturated or unsaturated 5-, 6- or 7-membered ring may optionally have one or more additional heteroatoms in the ring.

15 The term "heterocyclic" is used herein at all occurrences to mean a saturated or wholly or partially unsaturated 5-10-membered ring system (unless the cyclic ring system is otherwise limited) in which one or more rings contain one or more heteroatoms, including, but not limited to, pyrrolidine, piperidine, piperazine, morpholine, imidazolidine, pyrazolidine, and the like.

20 The term "CCR5 mediated disease state" is used herein at all occurrences to mean any disease state which is mediated (or modulated) by CCR5.

Suitably, pharmaceutically acceptable salts of formula (I) include, but are not limited to, salts with inorganic acids such as hydrochloride, sulfate, phosphate, diphosphate, hydrobromide, and nitrate, or salts with an organic acid such as malate, maleate, fumarate, tartrate, succinate, citrate, acetate, lactate, 25 methanesulfonate, p-toluenesulfonate, palmitate, salicylate, and stearate.

Among the preferred compounds of the invention are the following compounds:

N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-(1,1'-biphenyl)-4-carboxamide;

30 N-[3-(2-Methylamino)ethoxy-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-(1,1'-biphenyl)-4-carboxamide;

N-[3-(2-Amino)ethoxy-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-(1,1'-biphenyl)-4-carboxamide;

35 N-[3-(2-Dimethylamino-1-methyl)ethoxy-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-(1,1'-biphenyl)-4-carboxamide;

3-(2-Dimethylaminoethoxy)-4-methoxy-N-[2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)[1,1'-biphenyl]]-4-benzamide oxalate;

- N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-(1,1-biphenyl)-4-carboxamide oxalate;
- N-[3-(2-Dimethylaminoethoxy)-4-methoxyphenyl]-4'-(5-ethyl-1,2,4-oxadiazol-3-yl)-2'-methylbiphenyl-4-carboxamide;
- 5 N-[3-(2-Pyrrolidin-1-ylethoxy)-4-methoxyphenyl]-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-2'-methylbiphenyl-4-carboxamide;
- N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-2'-ethyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-(1,1'-biphenyl)-4-carboxamide hydrochloride;
- 10 N-(3-(2-Dimethylamino)ethoxy-4-methoxyphenyl)-2',3-dimethyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-1,1'-biphenyl-4-carboxamide hydrochloride;
- N-[3-((S)-1-Methylpyrrolidin-2-ylmethoxy)-4-methoxyphenyl]-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-2'-methyl-biphenyl-4-carboxamide;
- N-[3-(1-Methyl-3-piperidinyloxy)-4-methoxyphenyl]-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-2'-methyl-biphenyl-4-carboxamide oxalate;
- 15 N-[3-(1-Methyl-3-azepinyloxy)-4-methoxyphenyl]-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-2'-methyl-biphenyl-4-carboxamide oxalate;
- N-[4-Methoxy-3-(quinuclidine-3-yloxy)phenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carboxamide oxalate;
- N-[3-((R)-1-Methylpyrrolidin-2-ylmethoxy)-4-methoxyphenyl]-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-2'-methyl-biphenyl-4-carboxamide;
- 20 N-[3-(1-Methylpyrrolidin-3-yl)-4-methoxy-phenyl]-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-2'-methyl-biphenyl-4-carboxamide;
- N-[3-((R)-Pyrrolidin-2-ylmethoxy)-4-methoxyphenyl]-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-2'-methylbiphenyl-4-carboxamide hydrochloride;
- 25 N-[3-(Azetidin-2-ylmethoxy)-4-methoxyphenyl]-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-2'-methylbiphenyl-4-carboxamide hydrochloride;
- N-[2-(Dimethylaminomethyl)-1,4-benzodioxan-7-yl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carboxamide;
- N-[2-(dimethylaminomethyl)-1,4-benzodioxan-6-yl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carboxamide;
- 30 2,3-Dihydro-6-(2-dimethylaminoethoxy)-5-methoxy-1-[4-(2-methyl-4-(5-methyl-1,2,4-oxadiazol-3-yl)phenyl)benzoyl]indole;
- 6-(2-Dimethylaminoethoxy)-5-methoxy-1-[4-(2-methyl-4-(5-methyl-1,2,4-oxadiazol-3-yl)phenyl)benzoyl]indole;
- 35 2,3-Dihydro-6-(2-dimethylaminoethoxy)-5-methoxy-2-methyl-1-[4-(2-methyl-4-(5-methyl-1,2,4-oxadiazol-3-yl)phenyl)benzoyl]indole oxalate;

- 6-(2-Dimethylaminoethoxy)-5-methoxy-2-methyl-1-[4-(2-methyl-4-(5-methyl-1,2,4-oxadiazol-3-yl)phenyl)benzoyl]indole oxalate;
- [7-(2-Dimethylaminoethoxy)-6-methoxy-2-methyl-3,4-dihydro-2H-quinolin-1-yl][2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-biphenyl-4-yl]methanone oxalate;
- 5 [8-(2-Dimethylaminoethoxy)-7-methoxy-2,3,4,5-tetrahydro-1H-benz[b]azepin-1-yl]-[2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-biphenyl-4-yl]methanone oxalate;
- 5-[4-(2-Methyl-4-(5-methyl-1,2,4-oxadiazol-3-yl)phenyl)-benzoylamino]spiro[(2,3-dihydrobenzofuran)-3,4'-piperidine] hydrochloride;
- 10 2,3,6,7-Tetrahydro-1'-methyl-5-[2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carbonyl]furo[2,3-f]indole-3-spiro-4'-piperidine hydrochloride;
- 5-[4'-(5-Dimethylamino-1,2,4-oxadiazol-3-yl)-2'-methylbiphenyl-4-carbonyl]-1'-methyl-2,3,5,6,7,8-hexahydrospiro(furo[2,3-g]quinoline-3,4'-piperidine) hydrochloride;
- 15 5-[4'-(5-Hydroxymethyl-1,2,4-oxadiazol-3-yl)-2'-methylbiphenyl-4-carbonyl]-1'-methyl-2,3,6,7-tetrahydrospiro[furo[2,3-f]indole-3,4'-piperidine] hydrochloride;
- 5-[4'-(5-Methoxymethyl-1,2,4-oxadiazol-3-yl)-2'-methylbiphenyl-4-carbonyl]-1'-methyl-2,3,6,7-tetrahydrospiro[furo[2,3-f]indole]-3,4'-piperidine hydrochloride;
- 20 4'-ax-(Dimethylamino)-5-[2'-methyl-4'-[(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-yl]carbonyl]-2,3,6,7-tetrahydrospiro[furo[2,3-f]indole-3,1'-cyclohexane] hydrochloride.
- 25 N-[3-(2-Dimethylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[4-(2-Dimethylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[2-(3-Dimethylamino)propoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[3-[2-(Morpholin-1-yl)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;
- 30 N-[3-(2-Dicyclohexylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-Ethyl-N-[3-[2-(N-isopropyl-N-methylamino)ethoxy]-4-methoxyphenyl]-1,1'-biphenyl-4-carboxamide;
- N-Methyl-N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-(1,1-biphenyl)-4-carboxamide;
- 35

- N-Methyl-N-[3-(diisopropylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4-iodobenzamide;
- N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-3-bromobenzamide;
- 5 N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4-(methoxy)benzamide;
- N-[4-(2-Diisopropylamino)ethoxy-phenyl]-4-iodobenzamide;
- N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-3-iodobenzamide;
- N-[3-(2-Diisopropylamino)ethoxy-phenyl]-3-iodobenzamide;
- 10 N-[3-(2-Diethylamino)ethoxy-phenyl]-3-iodobenzamide;
- N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-4-bromobenzamide;
- N-[4-(2-Diisopropylamino)ethoxy-phenyl]-4-bromobenzamide;
- N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-3-bromobenzamide;
- N-[4-(2-Diisopropylamino)ethoxy-phenyl]-4-(isopropyl)benzamide;
- 15 N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-4-(cyclohexyl)benzamide;
- N-[4-(2-Diethylamino)ethoxy-phenyl]-4-(cyclohexyl)benzamide;
- N-[4-(2-Diisopropylamino)ethoxy-phenyl]-4'-ethyl-[1,1'-biphenyl]-4-carboxamide;
- 20 N-[4-(2-Diethylamino)ethoxy-phenyl]-4'-ethyl-[1,1'-biphenyl]-4-carboxamide;
- N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-4'-methoxy-[1,1'-biphenyl]-4-carboxamide;
- N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-4'-hydroxy-[1,1'-biphenyl]-4-carboxamide;
- 25 N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-3'-hydroxy-[1,1'-biphenyl]-4-carboxamide;
- N-[2-(dimethylaminomethyl)-1,4-benzodioxan-7-yl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carboxamide; and
- 30 N-[2-(dimethylaminomethyl)-1,4-benzodioxan-6-yl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carboxamide.

Among the more preferred compounds of the invention are the following compounds:

- N-[3-(3-Dimethylamino)propoxy-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-(1,1'-biphenyl)-4-carboxamide;
- 5 N-[3-(2-Piperidine)ethoxy-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-(1,1' biphenyl)-4-carboxamide;
- N-[3-(3-Dimethylaminopropyl)-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carboxamide oxalate;
- 10 N-[3-(1-Methyl-4-piperidyl)-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carboxamide oxalate;
- N-[3-(1-Methyl-3-piperidyl)methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carboxamide oxalate;
- N-[7-(2-Dimethylamino)ethoxy-2,3-dihydrobenzofuran-5-yl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-1,1'-biphenyl-4-carboxamide;
- 15 [3-(Dimethylaminoethyl)-3,6,7,8-tetrahydro-2H-furo[2,3-g]quinolin-5-yl]-[2'-methyl-4'-(5-methyl-[1,2,4]-oxadiazol-3-yl)-biphenyl-4-yl]-methanone oxalate;
- 5-[4-(2-Methyl-4-(5-methyl-1,2,4-oxadiazol-3-yl)phenyl)benzoylamino]-spiro[(2,3-dihydrobenzofuran)-3,4'-(1-methylhexahydroazepine)] oxalate;
- 2,3,5,6,7,8,-Hexahydro-1'-methyl-5-{2'-methyl-4'-[(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-yl]carbonyl} furo[2,3-g]quinoline-3-spiro-4'-piperidine
- 20 oxalate;
- 6,7,8,9-Tetrahydro-1'-methyl-5-[[2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)[1,1'-biphenyl]-4-yl]carbonyl]-spiro[2H-furo[2,3-h][1]benzazepine-3(5H),4'-piperidine];
- 25 2,3,5,6,7,8-Hexahydro-5-[4'-(5-hydroxymethyl-1,2,4-oxadiazol-3-yl)-2'-methylbiphenyl-4-carbonyl]-1'-methylspiro[furo[2,3-g]quinoline-3,4'-piperidine] hydrochloride;
- 4'-(Dimethylamino)-5-[2'-methyl-4'-[(5-methyl-1,2,4-oxadiazol-3-yl)-biphenyl-4-yl]carbonyl]-2,3,6,7-tetrahydrospiro[furo[2,3-f]indole-3,1'-cyclohexane] hydrochloride;
- 30 N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-3-carboxamide;
- N-[3-(3-Dimethylamino)propoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[4-(3-Dimethylamino)propoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide;
- 35 N-[3-(Diethylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;

- N-[3-[2-(Piperidin-1-yl)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[3-(2-Diisopropylamino)ethoxy-2-methylphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-[1,1'-biphenyl]-4-carboxamide;
- 5 N-[3-(2-Dipropylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[3-(2-Diisopropylamino)ethoxy-4-methylphenyl]-[1,1'-biphenyl]-4-carboxamide;
- 10 N-[3-(2-Diisopropylamino)ethoxy-4-ethylphenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[3-(2-Diisopropylamino)ethoxy-4-(methoxycarbonylamino)phenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[1,1'-Biphenyl]-4-yl-3-(2-diisopropylamino)ethoxy-4-methoxybenzamide;
- 15 N-[3-(2-Diisopropylamino)ethoxy-4-(ethoxycarbonyl)phenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[4-(2-Diisopropylamino)ethoxy-3-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;
- 20 N-[2-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[3-(2-diisopropylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[2-Fluoro-4-(2-diisopropylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide;
- 25 N-[4-(2-Diisopropylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[2-(2-Diisopropylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4-bromobenzamide;
- N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4-(isopropyl)benzamide;
- 30 N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4-(cyclohexyl)benzamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-3,4-dichlorobenzamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-3,5-dichlorobenzamide;

5 N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4-(nitro)benzamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-3-iodobenzamide;

N-[4-(2-Diisopropylamino)ethoxy-phenyl]-3-iodobenzamide;

N-[4-(2-Diisopropylamino)ethoxy-phenyl]-3-bromobenzamide;

N-[4-(2-Diisopropylamino)ethoxy-phenyl]-4-(cyclohexyl)benzamide;

10 N-[3-(2-Diisopropylamino)ethoxy-4-methoxy-phenyl]-4-(cyclohexyl)benzamide;

N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-4'-ethyl-[1,1'-biphenyl]-4-carboxamide;

15 N-[3-(2-Diisopropylamino)ethoxy-phenyl]-4'-ethyl-[1,1'-biphenyl]-4-carboxamide; and

N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-3'-methoxy-[1,1'-biphenyl]-4-carboxamide).

Among the most preferred compounds of the invention are the following compounds:

20 N-[3-(2-Diisopropylamino)ethoxy-phenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-(1,1'-biphenyl)-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-(1,1'-biphenyl)-4-carboxamide;

25 5-[4-(2-methyl-4-(5-methyl-1,2,4-oxadiazol-3-yl)phenyl)benzoylamino]spiro[(2,3-dihydrobenzofuran)-3,4'-(1-methylpiperidine)] oxalate;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-(1,1'-biphenyl)-4-carboxamide;

30 N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-2'-methyl-(1,1'-biphenyl)-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-phenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-[1,1'-biphenyl]-4-carboxamide;

- N-[3-[2-(N-Cyclohexyl-N-isopropylamino)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-carboxamide;
- N-[3-[2-(cis-2,6-Dimethylpiperidin-1-yl)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;
- 5 N-[3-[2-(N-Ethyl-N-isopropylamino)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[3-[2-(2,5-Dimethylpyrrolidin-1-yl)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[4-Methoxy-3-[2-(2,2,6,6-tetramethylpiperidin-1-yl)ethoxy]phenyl]-1,1'-biphenyl-4-carboxamide;
- 10 N-[3-[2-(N-t-Butyl-N-isopropyl)aminoethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[3-(3-Diisopropylamino)propyloxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;
- 15 N-[3-(3-Diisopropylamino)propyl-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[3-(2-Diisopropylamino)ethoxy-4-iodophenyl]-[1,1'-biphenyl]-4-carboxamide;
- N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-methanamine Dihydrochloride;
- 20 N-[3-(2-Diisopropylamino)ethoxy-4-methoxy-phenyl]-4'-cyano-1,1'-biphenyl-4-carboxamide;
- N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4'-cyano-2'-methyl-1,1'-biphenyl-4-carboxamide;
- 25 N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-2-naphthalenecarboxamide;
- N-(1'-Methylspiro[benzofuran-3(2H),4'-piperidin]-5-yl)-[1,1'-biphenyl]-4-carboxamide;
- N-[3-[2-(N-Isopropyl-N-methylamino)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;
- 30 N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4-(iodo)benzamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxy-phenyl]-4-(cyclohexyl)benzamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4'-ethyl-[1,1'-biphenyl]-4-carboxamide;

5 N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-3'-methoxy-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4'-acetyl-[1,1'-biphenyl]-4-carboxamide;

10 N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-3'-methoxy-[1,1'-biphenyl]-4-carboxamide; and

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-3'-hydroxy-[1,1'-biphenyl]-4-carboxamide.

Compounds excluded from the scope of this invention are as follows:

15 N-[3-(2-Dimethylaminoethoxy)-4-methoxyphenyl]-4'-(5-dimethylamino-1,2,4-oxadiazol-3-yl)-2'-methylbiphenyl-4-carboxamide;

N-[3-(1-Methylazetidin-2-ylmethoxy)-4-methoxyphenyl]-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-2'-methylbiphenyl-4-carboxamide;

[7-(2-Dimethylaminoethoxy)-6-methoxy-3,4-dihydro-2H-quinolin-1-yl]-[2'-methyl-[1,2,4]-oxadiazol-3-yl)biphenyl-4-yl]-methanone oxalate;

20 2,3,6,7-Tetrahydro-1'-methyl-5-[2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carbonyl]-furo[2,3-f]indole-3-spiro-3'-piperidine oxalate;

5-[4'-(5-Methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carbonyl]-1'-methyl-2,3,6,7-tetrahydrofuro[2,3-f]indole-3-spiro-4'-piperidine;

1'-Ethyl-5-[2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-biphenyl-4-carbonyl]-2,3,6,7-tetrahydrospirofuro[2,3-f]indole-3,4'-piperidine;

25 N-[2-(2-Dimethylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]benzamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-2-carboxamide;

30 N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-4-(methoxy)benzamide;

N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-4-(nitro)benzamide;

- N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-4-(nitro)benzamide;
N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-4-(nitro)benzamide;
N-[3-(Dimethylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-
carboxamide;
- 5 N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-4'-chloro-[1,1'-
biphenyl]-4-carboxamide;
N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-4'-hydroxy-[1,1'-
biphenyl]-4-carboxamide;
N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-4'-acetyl-[1,1'-biphenyl]-
10 4-carboxamide;
N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-4'-acetyl-[1,1'-
biphenyl]-4-carboxamide;
N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-3'-chloro-[1,1'-
biphenyl]-4-carboxamide;
- 15 N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-3'-nitro-[1,1'-biphenyl]-4-
carboxamide;
N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-3'-nitro-[1,1'-biphenyl]-
4-carboxamide;
- 20 N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-4-iodobenzamide;
N-[4-(2-Diethylamino)ethoxy-phenyl]-4-iodobenzamide;
N-[2-(2-Diisopropylamino)ethoxy-phenyl]-4-iodobenzamide;
N-[4-(2-Diethylamino)ethoxy-phenyl]-3-iodobenzamide;
N-[2-(2-Diisopropylamino)ethoxy-phenyl]-3-iodobenzamide;
N-[2-(2-Diethylamino)ethoxy-phenyl]-3-iodobenzamide;
- 25 N-[3-(2-Diethylamino)ethoxy-phenyl]-4-bromobenzamide;
N-[2-(2-Diisopropylamino)ethoxy-phenyl]-4-bromobenzamide;
N-[2-(2-Diethylamino)ethoxy-phenyl]-4-bromobenzamide;
N-[4-(2-Diethylamino)ethoxy-phenyl]-3-bromobenzamide;
N-[2-(2-Diisopropylamino)ethoxy-phenyl]-3-bromobenzamide;
- 30 N-[2-(2-Diethylamino)ethoxy-phenyl]-3-bromobenzamide;

- N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-4-(dimethylamino)benzamide;
- N-[4-(2-Diisopropylamino)ethoxy-phenyl]-4-(dimethylamino)benzamide;
- N-[4-(2-Diethylamino)ethoxy-phenyl]-4-(dimethylamino)benzamide;
- 5 N-[2-(2-Diisopropylamino)ethoxy-phenyl]-4-(dimethylamino)benzamide;
- N-[2-(2-Diethylamino)ethoxy-phenyl]-4-(dimethylamino)benzamide;
- N-[4-(2-Diethylamino)ethoxy-phenyl]-4-(isopropyl)benzamide;
- N-[2-(2-Diisopropylamino)ethoxy-phenyl]-4-(isopropyl)benzamide;
- N-[2-(2-Diethylamino)ethoxy-phenyl]-4-(isopropyl)benzamide;
- 10 N-[3-(2-Diethylamino)ethoxy-phenyl]-4-(cyclohexyl)benzamide;
- N-[2-(2-Diisopropylamino)ethoxy-phenyl]-4-(cyclohexyl)benzamide;
- N-[2-(2-Diethylamino)ethoxy-phenyl]-4-(cyclohexyl)benzamide;
- N-[2-(2-Diisopropylamino)ethoxy-phenyl]-4'-ethyl-[1,1'-biphenyl]-4-carboxamide; and
- 15 N-[2-(2-Diethylamino)ethoxy-phenyl]-4'-ethyl-[1,1'-biphenyl]-4-carboxamide.

Formulation of Pharmaceutical Compositions

The pharmaceutically effective compounds of this invention (and the pharmaceutically acceptable salts thereof) are administered in conventional dosage forms prepared by combining a compound of formula (I) ("active ingredient") in an amount sufficient to treat COPD, asthma and atopic disorders (for example, atopic dermatitis and allergies), rheumatoid arthritis, sarcoidosis and other fibrotic diseases, atherosclerosis, psoriasis, autoimmune diseases such as multiple sclerosis, inflammatory bowel disease, and HIV infection, ("CCR5-mediated disease states")

20 with standard pharmaceutical carriers or diluents according to conventional procedures well known in the art. These procedures may involve mixing, granulating and compressing or dissolving the ingredients as appropriate to the desired preparation.

The pharmaceutical carrier employed may be, for example, either a solid or liquid. Exemplary of solid carriers are lactose, terra alba, sucrose, talc, gelatin, agar, pectin, acacia, magnesium stearate, stearic acid and the like. Exemplary of liquid carriers are syrup, peanut oil, olive oil, water and the like. Similarly, the carrier or diluent may include time delay material well known to the art, such as glyceryl monostearate or glyceryl distearate alone or with a wax.

30

A wide variety of pharmaceutical forms can be employed. Thus, if a solid carrier is used, the preparation can be tableted, placed in a hard gelatin capsule in powder or pellet form or in the form of a troche or lozenge. The amount of solid carrier will vary widely but preferably will be from about 25 mg to about 1000 mg.

- 5 When a liquid carrier is used, the preparation will be in the form of a syrup, emulsion, soft gelatin capsule, sterile injectable liquid such as an ampule or nonaqueous liquid suspension.

- The active ingredient may also be administered topically to a mammal in need of treatment or prophylaxis of CCR5 mediated disease states. The amount of active ingredient required for therapeutic effect on topical administration will, of course, vary with the compound chosen, the nature and severity of the disease state being treated and the mammal undergoing treatment, and is ultimately at the discretion of the physician. A suitable dose of an active ingredient is 1.5 mg to 500 mg for topical administration, the most preferred dosage being 1 mg to 100 mg, for example 5 to 25 mg administered two or three times daily.

- By topical administration is meant non-systemic administration and includes the application of the active ingredient externally to the epidermis, to the buccal cavity and instillation of such a compound into the ear, eye and nose, and where the compound does not significantly enter the blood stream. By systemic administration is meant oral, intravenous, intraperitoneal and intramuscular administration.

- While it is possible for an active ingredient to be administered alone as the raw chemical, it is preferable to present it as a pharmaceutical formulation. The active ingredient may comprise, for topical administration, from 0.001% to 10% w/w, e.g. from 1% to 2% by weight of the formulation although it may comprise as much as 10% w/w but preferably not in excess of 5% w/w and more preferably from 0.1% to 1% w/w of the formulation.

- The topical formulations of the present invention, both for veterinary and for human medical use, comprise an active ingredient together with one or more acceptable carrier(s) therefor and optionally any other therapeutic ingredient(s). The carrier(s) must be 'acceptable' in the sense of being compatible with the other ingredients of the formulation and not deleterious to the recipient thereof.

- Formulations suitable for topical administration include liquid or semi-liquid preparations suitable for penetration through the skin to the site of inflammation such as liniments, lotions, creams, ointments or pastes, and drops suitable for administration to the eye, ear or nose.

Drops according to the present invention may comprise sterile aqueous or oily solutions or suspensions and may be prepared by dissolving the active ingredient in a

suitable aqueous or alcoholic solution of a bactericidal and/or fungicidal agent and/or any other suitable preservative, and preferably including a surface active agent. The resulting solution may then be clarified by filtration, transferred to a suitable container which is then sealed and sterilized by autoclaving or maintaining at 98-
5 100°C for half an hour. Alternatively, the solution may be sterilized by filtration and transferred to the container by an aseptic technique. Examples of bactericidal and fungicidal agents suitable for inclusion in the drops are phenylmercuric nitrate or acetate (0.002%), benzalkonium chloride (0.01%) and chlorhexidine acetate (0.01%). Suitable solvents for the preparation of an oily solution include glycerol, diluted
10 alcohol and propylene glycol.

Lotions according to the present invention include those suitable for application to the skin or eye. An eye lotion may comprise a sterile aqueous solution optionally containing a bactericide and may be prepared by methods similar to those for the preparation of drops. Lotions or liniments for application to the skin may also
15 include an agent to hasten drying and to cool the skin, such as an alcohol or acetone, and/or a moisturizer such as glycerol or an oil such as castor oil or arachis oil.

Creams, ointments or pastes according to the present invention are semi-solid formulations of the active ingredient for external application. They may be made by mixing the active ingredient in finely-divided or powdered form, alone or in solution
20 or suspension in an aqueous or non-aqueous fluid, with the aid of suitable machinery, with a greasy or non-greasy basis. The basis may comprise hydrocarbons such as hard, soft or liquid paraffin, glycerol, beeswax, a metallic soap; a mucilage; an oil of natural origin such as almond, corn, arachis, castor or olive oil; wool fat or its derivatives, or a fatty acid such as stearic or oleic acid together with an alcohol such
25 as propylene glycol. The formulation may incorporate any suitable surface active agent such as an anionic, cationic or non-ionic surfactant such as esters or polyoxyethylene derivatives thereof. Suspending agents such as natural gums, cellulose derivatives or inorganic materials such as siliceous silicas, and other ingredients such as lanolin, may also be included.

The active ingredient may also be administered by inhalation. By "inhalation" is meant intranasal and oral inhalation administration. Appropriate dosage forms for such administration, such as an aerosol formulation or a metered dose inhaler, may be prepared by conventional techniques. The daily dosage amount of the active
30 ingredient administered by inhalation is from about 0.1 mg to about 100 mg per day, preferably about 1 mg to about 10 mg per day.
35

In one aspect, this invention relates to a method of treating COPD, asthma and atopic disorders (for example, atopic dermatitis and allergies), rheumatoid

arthritis, sarcoidosis and other fibrotic diseases, atherosclerosis, psoriasis, autoimmune diseases such as multiple sclerosis, inflammatory bowel disease, and HIV infection, all in mammals, preferably humans, which comprises administering to such mammal an effective amount of a CCR5 receptor ligand, in particular, a
5 compound as depicted in formula (I).

By the term "treating" is meant either prophylactic or therapeutic therapy. Such formula (I) compound can be administered to such mammal in a conventional dosage form prepared by combining the formula (I) compound with a conventional pharmaceutically acceptable carrier or diluent according to known techniques. It
10 will be recognized by one of skill in the art that the form and character of the pharmaceutically acceptable carrier or diluent is dictated by the amount of active ingredient with which it is to be combined, the route of administration and other well-known variables. The formula (I) compound is administered to a mammal in need of treatment for asthma and atopic disorders (for example, atopic dermatitis
15 and allergies), rheumatoid arthritis, atherosclerosis, psoriasis, autoimmune diseases such as multiple sclerosis, inflammatory bowel disease, and HIV infection, in an amount sufficient to decrease symptoms associated with these disease states. The route of administration may be oral or parenteral.

The term parenteral as used herein includes intravenous, intramuscular, subcutaneous, intra-rectal, intravaginal or intraperitoneal administration. The
20 subcutaneous and intramuscular forms of parenteral administration are generally preferred. The daily parenteral dosage regimen will preferably be from about 30 mg to about 300 mg per day of active ingredient. The daily oral dosage regimen will preferably be from about 100 mg to about 2000 mg per day of active
25 ingredient.

It will be recognized by one of skill in the art that the optimal quantity and spacing of individual dosages of a formula (I) compound will be determined by the nature and extent of the condition being treated, the form, route and site of administration, and the particular mammal being treated, and that such optimums
30 can be determined by conventional techniques. It will also be appreciated by one of skill in the art that the optimal course of treatment, i.e., the number of doses of the formula (I) compound given per day for a defined number of days, can be ascertained by those skilled in the art using conventional course of treatment determination tests.

35 **Methods of Preparation**

The compounds of formula (I) can be prepared by art-recognized procedures from known or commercially available starting materials. If the

starting materials are unavailable from a commercial source, their synthesis is described herein, or they can be prepared by procedures known in the art.

Specifically, compounds of formula (I) wherein R is represented by group (a), A is CONH and E is represented by group (d), were prepared according to the methods of international application publication number WO 95/15954, published 15 June 1995.

Compounds of formula (I) wherein R is represented by group (a), A is CONH and E is represented by group (e), were prepared according to the methods of international application publication number WO 95/17401, published 29 June 1995.

Compounds of formula (I) wherein R is represented by group (a), A is CONH and E is represented by group (f), were prepared according the methods of international application publication number WO 96/31508 published 10 October 1996.

Compounds of formula (I) wherein R is represented by group (a), A is CONH and E is represented by groups (g) or (h), were prepared according the methods of international application publication number WO 95/30675, published 16 November 1995.

Compounds of formula (I) wherein R is represented by group (a), A is CONH and E is represented by group (j), were prepared according the methods of international application publication number WO 96/11934, published 25 April 1996.

Compounds of formula (I) wherein R is represented by group (a) and AE is represented by group (k), were prepared according the methods of international application publication number WO 95/17398, published 29 June 1995.

Compounds of formula (I) wherein R is represented by group (a) and AE is represented by group (l), were prepared according the methods of international application publication number WO 97/07120, published 27 February 1997.

Compounds of formula (I) wherein R is represented by group (a) and AE is represented by group (m), were prepared according the methods of international application publication number WO 96/19477, published 27 June 1996.

Compounds of formula (I) wherein R is represented by group (a) and AE is represented by group (n), were prepared according the methods of international application publication number WO 97/19070 published 29 May 1997.

Specifically, compounds of formula (I) wherein Ar is represented by group (i), (ii) or (iii), A is CONR⁴⁶, NHCO or CH₂NH, and E is represented by group (a), were prepared according to the methods of international application publication

number WO 95/15954, published 15 June 1995, international application publication number WO 95/17398, published 29 June 1995, international application publication number WO 95/263284, published 5 October 1995, international application publication number WO 96/06079, published 29 February 5 1996, GB 2276161 published 21 September 1994, and GB 2276165 published 21 September 1994.

Compounds of formula (I) wherein Ar is (i) or (ii), and A is CONR⁶⁰ or NHCO, and E is represented by group (b), were prepared according to the methods of international application publication number WO 95/11934, published 25 April 10 1995, and WO 95/19477, published 27 June 1995. Four other applications cover the spiro compounds WO 97/17350 published 15 May 1997; WO 97/34900 published 25 September 1997; WO 97/34901 published 25 September 1997; WO 97/35862 published 2 October 1997.

Compounds of formula (I) wherein Ar is (i), (ii) or (iii), A is CONR⁷⁰, 15 NHCO or CH₂NH, and E represents (c), were prepared according the methods of international application publication number WO 95/306758 published 16 November 1995 and GB 2276165 published 21 September 1994.

Compounds of formula (I) wherein Ar is (ii), A is CONH or NHCO, and E represents (f), were prepared according the methods of international application 20 publication number WO 95/17401, published 29 June 1995.

Compounds of formula (I) wherein Ar is (i) or (ii), A is CONR⁹³, and E represents a group (g), were prepared according the methods of international application publication number WO 96/31508, published 10 October 1996.

Compounds of formula (I) when Ar is (i) or (ii), A is CONR¹⁰¹, and E 25 represents group (h), were prepared according the methods of international application publication number WO 95/32767, published 7 December 1995 and WO 97/07120, published 27 February 1997.

Compounds of formula (I) Ar is (i) or (ii), and A is CONR¹⁰⁹ or CH₂NH, and E represents group (i), were prepared according the methods of international 30 application publication number WO 97/19070, published 29 May 1997.


Compounds of formula (I) are also prepared by the method of Scheme 1 using solid-phase chemistry. Resin 1-1 (WO 98/17695) is coupled with a suitably substituted carboxyphenyl-boronic acid, for Example 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoic acid, using Suzuki conditions, for example, a suitable 35 palladium reagent, such as tetrakis(triphenylphosphine)palladium(0), and a suitable base, such as sodium carbonate, in a suitable solvent, such as ethanol and toluene, at a suitable temperature, such as 90°C, for a suitable time, such as 24 h, to afford

the resin-bound biphenylcarboxylic acid **1-2**. The adduct **1-2** is condensed with appropriately substituted anilines **1-4**, prepared by methods known to the art from commercially available starting materials or as described herein, using methods known to the art. For example, treatment of **1-2** with a suitable reagent, such as

5 oxalyl chloride, in a suitable solvent, such as toluene, affords the biphenylcarbonyl chloride adduct **1-3**. Treatment of **1-3** with **1-4** in a suitable solvent, such as dichloromethane, in the presence of a suitable base, such as diisopropylethylamine, gives the biphenylcarboxanilide adduct, **1-5**. Treatment of **1-5** with a suitable acid and solvent, such as trifluoroacetic/dichloromethane/water (50:48:2) affords **1-6**

10 which are compounds of formula (I).


Scheme 1



a) 4-carboxybenzeneboronic acid, $(\text{Ph}_3\text{P})_4\text{Pd}$, Na_2CO_3 , EtOH, toluene, 90°C , 24 h; b) $(\text{COCl})_2$, toluene; c) 4, diisopropylethylamine, CH_2Cl_2 ; d) $\text{CF}_3\text{CO}_2\text{H}$, CH_2Cl_2 .

Compounds of formula (I) are also prepared by the method of Scheme 2 using solid-phase chemistry. Appropriately substituted (alkylamino)ethoxy-anilines **2-1**, such as 3-(2-diisopropylamino)-ethoxy-4-methoxyaniline, synthesized according to literature methods from commercially available starting materials and by methods described herein, are reacted with 4-formyl-3,5-dimethoxyphenol-Merrifield resin **2-2** (Boojamra, et al., *J. Org. Chem.* **1995**, 60, 5742) is treated and a suitable reducing agent, such as sodium triacetoxyborohydride, in a suitable solvent, such as dimethylformamide containing 1% acetic acid, to afford **2-3**. The resin-bound aniline **2-3** is condensed with an appropriately substituted benzoic acid or biphenylcarboxylic acid, which are commercially available or synthesized by methods known to the art, using a suitable activating agent, such as N-bromosuccinimide and triphenylphosphine, in a suitable solvent, such as dichloromethane, dimethylformamide and pyridine, to afford **2-4**. Treatment of **2-4** with a suitable acid and solvent, such as trifluoroacetic:dichloromethane:water (50:48:2) gives the desired benzanilide or biphenylcarboxanilide **2-5** which are compounds of formula (I).

Scheme 2



a) BAL resin, $\text{NaBH}(\text{OAc})_3$, 1% HOAc, DMF; b) X-benzoic acid, N-bromo-succinimide, Ph_3P , pyridine; c) TFA, CH_2Cl_2 , H_2O .

The invention will now be described by reference to the following examples which are merely illustrative and are not to be construed as a limitation of the scope of the present invention. In the Examples, mass spectra were performed upon a VG Zab mass spectrometer using fast atom bombardment, unless otherwise indicated.

EXAMPLES

10

Preparation 1

Preparation of 4-(2-Diisopropylamino)ethoxy-3-methoxyaniline

a) 4-(2-diisopropylamino)ethoxy-3-methoxy-1-nitrobenzene

A mixture of 4-nitroguaiacole (1.0 g, 5.9 mmol), 2-diisopropylaminoethyl chloride hydrochloride (1.2 g, 5.9 mmol), and powdered potassium carbonate (1.38 g, 10 mmol) in dry acetone (100 mL) was heated to reflux, stirred for 16 h, cooled, filtered, and the filtrate was concentrated *in vacuo*. The residue was partitioned between ethyl acetate (300 mL) and water (40 mL), and the organic phase was washed with water (40 mL), dried (Na₂SO₄), and concentrated *in vacuo* to give the title compound: MS(ES) m/e 297.2 [M+H]⁺.

b) 4-(2-diisopropylamino)ethoxy-3-methoxyaniline

A mixture of the compound of Preparation 1(a) (1.6 g, 5.4 mmol) and 5% palladium-on-carbon (0.3 g) in absolute ethanol (100 mL) was shaken in a hydrogen atmosphere (50 psi) for 2.5 h, filtered, and the filtrate was concentrated *in vacuo* to give the title compound: MS(ES) m/e 267.3 [M+H]⁺.

Preparations 2-8

Preparation of 2-(2-Diisopropylamino)ethoxy-4-methoxyaniline, 3-(2-Diisopropylamino)ethoxyaniline, 4-(2-Diisopropylamino)ethoxy-2-fluoroaniline, 4-(2-Diisopropylamino)ethoxyaniline, 2-(2-Diisopropylamino)ethoxyaniline, 3-(2-Diisopropylamino)ethoxy-2-methylaniline, and 3-(2-Diisopropylamino)ethoxy-4-methylaniline

Following the procedure of Preparation 1, except substituting 5-methoxy-2-nitrophenol, 3-nitrophenol, 3-fluoro-4-nitrophenol, 4-nitrophenol, 2-nitrophenol, 2-methyl-3-nitrophenol, or 2-methyl-5-nitrophenol for 4-nitroguaiacole, gave the title compounds:

2-(2-diisopropylamino)ethoxy-4-methoxyaniline: MS(ES) m/e 267.2 [M+H]⁺;
3-(2-diisopropylamino)ethoxyaniline: MS(ES) m/e 237.2 [M+H]⁺;
4-(2-diisopropylamino)ethoxy-2-fluoroaniline: MS(ES) m/e 255.0 [M+H]⁺;
4-(2-diisopropylamino)ethoxyaniline: MS(ES) m/e 237.2 [M+H]⁺;
2-(2-diisopropylamino)ethoxyaniline: MS(ES) m/e 237.1 [M+H]⁺;
3-(2-diisopropylamino)ethoxy-2-methylaniline: MS(ES) m/e 251.2 [M+H]⁺; and
3-(2-diisopropylamino)ethoxy-4-methylaniline: MS(ES) m/e 251.1 [M+H]⁺.

Preparation 9

Preparation of 3-(2-Dicyclohexylamino)ethoxy-4-methoxyaniline

a) N,N-dicyclohexyl-2-methoxy-5-nitro-phenoxyacetamide

A solution of 2-methoxy-5-nitro-phenoxyacetyl chloride (0.22 g, 0.9 mmol), prepared from 2-methoxy-5-nitro-phenoxyacetic acid (Brown et al., *J. Chem. Soc.* 1955, 3681) and thionyl chloride, in dichloromethane (10 mL) was added to a mixture of dicyclohexylamine (0.16 g, 0.9 mmol) and diisopropylethylamine (0.23 g, 1.76 mmol) in dichloromethane (20 mL), maintained at RT for 16 h, washed with water (2 x 20 mL), dried (Na₂SO₄), and concentrated *in vacuo*. The residue was recrystallized from methanol to afford the title compound (0.16 g).

b) N,N-dicyclohexyl-2-methoxy-5-amino-phenoxyacetamide
Following the procedure of Preparation 1(b), except substituting the compound of Preparation 9(a) for the compound of Preparation 1(a), gave the title compound.

c) 3-(2-dicyclohexylamino)ethoxy-4-methoxyaniline
A solution of the compound of Preparation 9(b) (0.15 g, 0.04 mmol) in anhydrous tetrahydrofuran (25 mL) was treated with 1.0M borane in tetrahydrofuran (4.3 mL, 4.3 mmol), heated to reflux for 3 h, maintained at RT for 16 h, carefully treated with methanol (5 mL), heated to reflux for 30 min, cooled, and concentrated *in vacuo*. The residue was partitioned between ethyl acetate (75 mL) and water (15 mL), and the organic phase was washed with water (15 mL), dried (Na₂SO₄), concentrated *in vacuo*, and the residue was purified by preparative TLC (Whatman PLK5F, 10% ethanol/dichloromethane-1% triethylamine) to give the title compound.

Preparation 10

Preparation 3-(2-Dipropylamino)ethoxy-4-methoxyaniline

a) 3-(2-dipropylamino)ethoxy-4-methoxy-1-nitrobenzene

A solution of 3-(2-bromoethoxy)-4-methoxy-1-nitrobenzene (Mutai et al., *Tetrahedron*, 1984, 40, 1755) (0.5 g, 1.8 mmol) and dipropylamine (1.8 g, 18 mmol) in dimethylformamide (10 mL) containing potassium iodide (0.28 g, 1.9 mmol) and potassium carbonate (0.42 g, 3 mmol) was warmed to 100°C and stirred for 16 h. The mixture was cooled, diluted with dichloromethane (100 mL), filtered, and the filtrate was concentrated *in vacuo*. The residue was partitioned between ethyl acetate (150 mL) and water (30 mL), and the organic phase was washed with water (4 x 30 mL), dried (Na₂SO₄), and concentrated *in vacuo* to give the title compound: MS(ES) m/e 297.1 [M+H]⁺.

b) 3-(2-dipropylamino)ethoxy-4-methoxyaniline

Following the procedure of Preparation 1(b), except substituting the compound of Preparation 10(a) for the compound of Preparation 1(a), gave the title compound: MS(ES) m/e 267.0 $[M+H]^+$.

5

Preparations 11-16

Preparation of 3-[2-(N-Cyclohexyl-N-isopropylamino)ethoxy]-4-methoxyaniline, 3-[2-(cis-2,6-Dimethylpiperidin-1-yl)ethoxy]-4-methoxyaniline, 3-[2-(N-Ethyl-N-isopropylamino)ethoxy]-4-methoxyaniline, 3-[2-(2,5-Dimethylpyrrolidin-1-yl)ethoxy]-4-methoxyaniline, 4-Methoxy-3-[2-(2,2,6,6-tetramethylpiperidin-1-yl)ethoxy]aniline, and 3-[2-(N-t-Butyl-N-isopropyl)aminoethoxy]-4-methoxyaniline

Following the procedure of Preparation 10, except substituting N-isopropylcyclohexylamine, cis-2,6-dimethylpiperidine, N-ethylisopropylamine, 2,5-dimethylpyrrolidine, 2,2,6,6-tetramethylpiperidine, or N-tert-butylisopropylamine for dipropylamine, gave the title compounds:

3-[2-(N-cyclohexyl-N-isopropylamino)ethoxy]-4-methoxyaniline: MS(ES) m/e 307.2 $[M+H]^+$;

3-[2-(cis-2,6-dimethylpiperidin-1-yl)ethoxy]-4-methoxyaniline: MS(ES) m/e 279.3 $[M+H]^+$;

3-[2-(N-ethyl-N-isopropylamino)ethoxy]-4-methoxyaniline: MS(ES) m/e 253.1 $[M+H]^+$;

3-[2-(2,5-dimethylpyrrolidin-1-yl)ethoxy]-4-methoxyaniline: MS(ES) m/e 265.1 $[M+H]^+$;

4-methoxy-3-[2-(2,2,6,6-tetramethylpiperidin-1-yl)ethoxy]aniline: MS(ES) m/e 307.0 $[M+H]^+$; and

3-[2-(N-t-butyl-N-isopropyl)aminoethoxy]-4-methoxyaniline: MS(ES) m/e 281 $[M+H]^+$.

Preparations 17-18

Preparation of 3-[2-(N-Isopropyl-N-methylamino)ethoxy]-4-methoxyaniline and N-Ethyl-3-[2-(N-isopropyl-N-methylamino)ethoxy]-4-methoxyaniline

a) 3-[2-(N-isopropyl-N-methylamino)ethoxy]-4-methoxy-1-nitrobenzene

Following the procedure of Preparation 10(a), except substituting N-methylisopropylamine for dipropylamine, gave the title compound: MS(ES) m/e 269.4 $[M+H]^+$.

b) 3-[2-(N-isopropyl-N-methylamino)ethoxy]-4-methoxyaniline and N-ethyl-3-[2-(N-isopropyl-N-methylamino)ethoxy]-4-methoxyaniline

Following the procedure of Preparation 1(b), except substituting the compound of Preparation 17(a) for the compound of Preparation 1(a), gave a mixture of the title compounds:

3-[2-(N-isopropyl-N-methylamino)ethoxy]-4-methoxyaniline: MS(ES) m/e
5 239.1 [M+H]⁺; and

N-ethyl-3-[2-(N-isopropyl-N-methylamino)ethoxy]-4-methoxyaniline:
MS(ES) m/e 267.1 [M+H]⁺.

Preparation 19

10 Preparation of 3-(3-Diisopropylamino)propyloxy-4-methoxyaniline

a) 3-(3-bromopropoxy)-4-methoxy-1-nitrobenzene

A stirred mixture of 2-methoxy-5-nitrophenol (5.0 g, 29.6 mmol) and powdered potassium carbonate (6.5 g, 47.4 mmol) in dimethylformamide (50 mL) was treated with 1,3-dibromopropane (80.8 g, 0.4 mol), heated to 90°C for 2 h,
15 cooled, diluted with dichloromethane (250 mL), and filtered. The filtrate was concentrated *in vacuo* and the residue was crystallized from methanol. The resulting solid was triturated with 30% ethyl acetate/hexane and then with boiling 30% ethyl acetate/hexane. The combined organic phases were concentrated *in vacuo* to give the title compound: ¹H NMR (300 MHz, CDCl₃) δ 7.92 (dd, 1H),
20 7.75 (d, 1H), 6.92 (dd, 1H), 4.20 (t, 2H), 3.95 (s, 3H), 3.62 (t, 2H), 2.40 (m, 2H).

b) 3-(3-diisopropylamino)propyloxy-4-methoxyaniline

Following the procedure of Preparation 1(b), except substituting the compound of Preparation 19(a) for the compound of Preparation 1(a), gave the title compound: MS(ES) m/e 281.1 [M+H]⁺.

25

Preparation 20

Preparation of 3-(3-Diisopropylamino)propyl-4-methoxyaniline

a) N,N-diisopropyl-3-(2-methoxy-5-nitrophenyl)propionamide

A solution of 3-(2-methoxy-5-nitrophenyl)propionyl chloride (0.46 g, 1.9
30 mmol), prepared from 3-(2-methoxy-5-nitrophenyl)propionic acid (Asano et al., *J. Pharm. Soc. Japan*, 1950, 70, 480) and thionyl chloride, in dichloromethane (5 mL) was added in one portion to a solution of diisopropylamine (0.67 g, 6.6 mmol) in dichloromethane (20 mL), stirred for 16 h, washed with water (2 x 20 mL), dried (Na₂SO₄), and concentrated *in vacuo* to give the title compound: MS(ES) m/z
35 309.1 [M+H]⁺.

b) N,N-diisopropyl-3-(2-methoxy-5-nitrophenyl)propylamine

A solution of the compound of Preparation 20(a) (0.6 g, 1.95 mmol) in dry tetrahydrofuran (50 mL) was treated with 1.0M borane in tetrahydrofuran (15 mL, 15 mmol), heated to reflux for 2 h, allowed to cool, stirred for 16 h, and carefully treated with methanol (10 mL). The mixture was concentrated *in vacuo*, and the residue was dissolved in dichloromethane (250 mL) and washed with water (2 x 40 mL). The organic phase was dried (Na₂SO₄) and concentrated *in vacuo* to give the title compound (0.52 g): MS(ES) m/z 295.1 [M+H]⁺.

c) 3-(3-diisopropylamino)propyl-4-methoxyaniline

Following the procedure of Preparation 1(b), except substituting the compound of Preparation 20(b) for the compound of Preparation 1(a), gave the title compound: MS(ES) m/e 265.3 [M+H]⁺.

Preparation 21

Preparation of 3-(2-Diisopropylamino)ethoxy-4-iodoaniline

a) 3-(2-diisopropylamino)ethoxy-4-iodo-1-nitrobenzene

Following the procedure of Preparation 1(a), except substituting 2-iodo-5-nitrophenol (Cambie et al., *Australian J. Chem.*, 1997, 50, 767) for 4-nitroguaiacole, gave the title compound: MS(ES) m/e 392.5 [M+H]⁺.

b) 3-(2-diisopropylamino)ethoxy-4-iodoaniline

A mixture of the compound of Preparation 21(a) (0.5 g, 1.27 mmol) and ferrous sulfate heptahydrate (10.0 g, 36 mmol) in water (50 mL) was stirred, heated to 100°C, and treated with concentrated ammonium hydroxide (6 mL) added dropwise over 5 min. The mixture was stirred for 5 min, cooled, filtered, and the filtrate was extracted with ethyl acetate. The organic phase was dried, concentrated *in vacuo*, and the residue was chromatographed (silica gel, 5% methanol/dichloromethane) to afford the title compound as a yellow solid (0.37 g): MS(ES) m/e 362.8 [M+H]⁺.

Preparation 22

Preparation of 3-(2-Diisopropylamino)ethoxy-4-ethylaniline Hydrochloride

a) 3-(2-diisopropylamino)ethoxy-4-(trimethylsilylethynyl)-1-nitrobenzene

A mixture of the compound of Preparation 21(a) (2.0 g, 5 mmol), trimethylsilylacetylene (0.8 g, 8 mmol), palladium acetate (15 mg), and triphenylphosphine (30 mg) in degassed triethylamine (25 mL) was heated to 90°C for 16 h. The mixture was concentrated *in vacuo* and the residue was partitioned between ethyl acetate and water. The organic phase was washed with brine, dried, and concentrated *in vacuo*. The residue was purified by flash chromatography

(silica gel, step gradient, 0-1% methanol/dichloromethane-ammonia) to afford the title compound (1.0 g): MS(ES) m/e 363 [M+H]⁺.

b) 3-(2-diisopropylamino)ethoxy-4-ethynyl-1-nitrobenzene

A solution of the compound of Preparation 22(a) (1 g, 2.7 mmol) in
5 methanol (20 mL) was treated with potassium carbonate (100 mg), stirred for 5 h, concentrated *in vacuo*, and the residue was dissolved in dichloromethane, and washed with 5% aqueous sodium carbonate and with brine. The organic phase was dried (MgSO₄) and concentrated *in vacuo* to afford the title compound (0.75 g): MS(ES) m/e 291.1 [M+H]⁺.

10 c) 3-(2-diisopropylamino)ethoxy-4-ethylaniline hydrochloride

A mixture of the compound of Preparation 22(b) (0.73 g, 2.5 mmol) (0.6 g) and 10% palladium-on-carbon (200 mg) in 1:1 methanol/ethanol (60 mL) was shaken in a hydrogen atmosphere for 4 h, filtered through Supercel, and the filtrate was concentrated *in vacuo*. The residue was dissolved in ethyl acetate, treated with
15 ethereal hydrogen chloride, stirred, and filtered to give the title compound (0.6 g): MS(ES) m/e 265.1 [M+H]⁺.

Preparation 23

Preparation of 3-(2-Diisopropylamino)ethoxy-4-(methoxycarbonylamino)aniline

20 a) 3-(2-diisopropylamino)ethoxy-4-nitroaniline hydrochloride

Following the procedure of Preparation 1(a), except substituting 2-amino-5-nitrophenol for 4-nitroguaiacole, gave the title compound: MS(ES) m/e 281.9 [M+H]⁺.

25 b) 3-(2-diisopropylamino)ethoxy-4-(methoxycarbonylamino)-1-nitrobenzene

A solution of the compound of Preparation 23(a) (1.27 g, 4 mmol) and diisopropylethylamine (2.1 mL, 12 mmol) in dichloromethane (25 mL) was treated with methyl chloroformate (0.38 g, 4 mmol), stirred for 7 d, treated with water, and the organic phase was washed with brine, dried, and concentrated *in vacuo*. The
30 residue was purified by flash chromatography (silica gel, step gradient, 0-1% methanol/dichloromethane-ammonia) to afford the title compound: MS(ES) m/e 339.9 [M+H]⁺.

c) 3-(2-diisopropylamino)ethoxy-4-(methoxycarbonylamino)aniline

Following the procedure of Preparation 1(b), except substituting the
35 compound of Preparation 23(b) for the compound of Preparation 1(a), gave the title compound: MS(ES) m/e 310.0 [M+H]⁺.

Preparation 24

Preparation of 3-(2-Diisopropylamino)ethoxy-4-methoxy-N-methyl-aniline

a) N-formyl-3-(2-diisopropylamino)ethoxy-4-methoxyaniline

A mixture of formic acid (0.3 g, 6.4 mmol) and acetic anhydride (0.53 g, 5.2 mmol) was heated to 50°C for 2 h, cooled, diluted with anhydrous tetrahydrofuran (10 mL), and treated in one portion with a solution of 3-(2-diisopropylamino)ethoxy-4-methoxyaniline (WO 95/15954) (0.5 g, 2 mmol) in tetrahydrofuran (5 mL). The resulting mixture was stirred for 16 h, concentrated *in vacuo*, and the residue was partitioned between ethyl acetate (85 mL) and 5% aqueous sodium carbonate (10 mL). The organic phase was washed with 5% sodium carbonate (10 mL) and then with brine (10 mL), dried (Na₂SO₄), and concentrated *in vacuo* to afford the title compound (0.57 g): MS(ES) m/e 295.4 [M+H]⁺.

b) 3-(2-diisopropylamino)ethoxy-4-methoxy-N-methyl-aniline

A solution of the compound of Preparation 24(a) (0.57 g, 1.9 mmol) in dry tetrahydrofuran (25 mL) was treated with 2.0M borane in tetrahydrofuran (2.4 mL, 4.8 mmol), stirred, and heated to reflux for 3 h. The mixture was cooled, maintained at RT for 18 h, carefully treated with methanol (10 mL), heated to reflux for 1 h, cooled, and concentrated *in vacuo*. The residue was partitioned between ethyl acetate (25 mL) and water (25 mL), and the organic phase was washed with water (25 mL) and with brine (25 mL), dried (Na₂SO₄), and concentrated *in vacuo* to afford the title compound (0.5 g): MS(ES) m/e 281.4 [M+H]⁺.

Preparations 25-26

Preparation of 2-Dimethylaminomethyl-1,4-benzodioxan-6-amine and 2-Dimethylaminomethyl-1,4-benzodioxan-7-amine

a) 2-chloromethyl-6-nitro-1,4-benzodioxan and 2-chloromethyl-7-nitro-1,4-benzodioxan

Following the procedure of *Gazz. Chim. Ital.*, 87, 1038-49 (1958), 2-chloromethyl-1,4-benzodioxan (2 g, 11 mmol) in glacial acetic acid (15 mL) was stirred, cooled, and treated with nitric acid (11 mL) added over 5 min. The mixture was heated to 95°C for 40 min, cooled, and carefully poured into water. The resulting mixture was extracted with chloroform, and the organic layer was washed and dried (Na₂SO₄). The residue was distilled, bp 166-8°C/1 mm Hg, to give a mixture of the title compounds (1.39 g).

b) 2-dimethylaminomethyl-6-nitro-1,4-benzodioxan and 2-dimethylaminomethyl-7-nitro-1,4-benzodioxan

The compounds of Preparation 25(a)-26(a)(1.64 g, 7 mmol) were dissolved in excess dimethylamine (15 mL) and heated to 150°C in a pressure vessel for 16

- 5 h. The mixture was cooled, triturated with ether, and filtered. The filtrate was concentrated *in vacuo* and the residue was distilled, bp 120-30°C/1 mm Hg, to give the title compounds (788 mg) as an orange oil. The product was purified further by acid/base treatment to give a mixture of the title compounds (600 mg).

- c) 2-dimethylaminomethyl-1,4-benzodioxan-6-amine and 2-dimethylaminomethyl-1,4-benzodioxan-7-amine

- The compounds of Preparation 25(b)-26(b)(600 mg, 2.5 mmol) and 10% palladium-on-carbon (200 mg) in ethanol (30 mL) were stirred under hydrogen (1 atm) for 1 h. The mixture was filtered, concentrated *in vacuo*, and chromatographed (silica gel, step gradient, 0-4% methanol/chloroform) followed by thick layer chromatography (step gradient, 1-2% methanol/chloroform) to give the title compounds:

2-dimethylaminomethyl-1,4-benzodioxan-6-amine, TLC R_f 0.56 (silica gel, 10% ethanol/chloroform); and

- 2-dimethylaminomethyl-1,4-benzodioxan-7-amine, TLC R_f 0.51 (silica gel, 10% ethanol/chloroform).

Example 1

Preparation of N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide

- 25 A solution of 4-biphenylcarbonyl chloride (0.32 g, 1.5 mmol), prepared from 4-biphenylcarboxylic acid and thionyl chloride, was added in one portion to a solution 3-(2-diisopropylamino)ethoxy-4-methoxyaniline (WO 95/15954)(0.37 g, 1.5 mmol) and diisopropylethylamine (0.19 g, 1.5 mmol) in dichloromethane (20 mL). The resulting mixture was stirred for 16 h, extracted with 5% aqueous sodium carbonate (10 mL), dried (Na₂SO₄), and concentrated *in vacuo*. The residue was purified by flash chromatography (silica gel, 10% methanol/dichloromethane) to give the title compound (0.22 g): MS(ES) m/e 447.1 [M+H]⁺.

35

Example 2

Preparation of N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-2'-methyl-[1,1'-biphenyl]-4-carboxamide

Following the procedure of Example 1, except substituting 2'-methyl-4-biphenylcarboxylic acid (Klein et al., *J. Med. Chem.* 1998, 41, 437) for 4-biphenylcarboxylic acid, gave the title compound: MS(ES) m/e 461.3 [M+H]⁺.

5

Example 3

Preparation of N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-3-carboxamide

Following the procedure of Example 1, except substituting 3-biphenylcarboxylic acid for 4-biphenylcarboxylic acid, gave the title compound:
10 MS(ES) m/e 447.3 [M+H]⁺.

Examples 4-11

Preparation of N-[3-(2-Dimethylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide; N-[4-(2-Dimethylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide; N-[2-(3-Dimethylamino)propoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide; N-[3-(3-Dimethylamino)propoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide; N-[4-(3-Dimethylamino)propoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide; N-[3-(Diethylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide; N-[3-[2-(Piperidin-1-yl)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide; and N-[3-[2-(4-Morpholinyl)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide
15
20

Following the procedure of Example 1, except substituting 3-(2-dimethylamino)ethoxyaniline (WO 95/26328), 4-(2-dimethylamino)ethoxyaniline (WO 96/23769), 2-(3-dimethylamino)-propoxyaniline (WO/9515954), 3-(3-dimethylamino)propoxyaniline (WO95/15954), 4-(3-dimethylamino)propoxyaniline (US 3,994,900), 3-(2-diethylamino)ethoxy-4-methoxyaniline (WO 95/15954), 3-[2-(piperidin-1-yl)ethoxy]-4-methoxyaniline (WO 95/15954), or 3-[2-(4-morpholinyl)ethoxy]-4-methoxyaniline (WO 95/15954) for 3-(2-diisopropylamino)ethoxy-4-methoxyaniline, gave the title compounds:

30 N-[3-(2-dimethylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide:
MS(ES) m/e 361 [M+H]⁺;

N-[4-(2-dimethylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide:
MS(ES) m/e 361 [M+H]⁺;

N-[2-(3-dimethylamino)propoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide:
35 MS(ES) m/e 375 [M+H]⁺;

N-[3-(3-dimethylamino)propoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide:
MS(ES) m/e 373 [M-H]⁺;

N-[4-(3-dimethylamino)propoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide:
MS(ES) m/e 375 [M+H]⁺;

N-[3-(diethylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-
carboxamide: MS(ES) m/e 419.5 [M+H]⁺;

5 N-[3-[2-(piperidin-1-yl)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-
carboxamide: MS(ES) m/e 431.3 [M+H]⁺; and

N-[3-[2-(4-morpholinyl)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-
carboxamide: MS(ES) m/e 433.0 [M+H]⁺.

10

Examples 12-13

Preparation of N-[3-(2-Diisopropylamino)ethoxy-2-methylphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-[1,1'-biphenyl]-4-carboxamide and N-[3-(2-Diisopropylamino)ethoxy-phenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-[1,1'-biphenyl]-4-carboxamide

15 Following the procedure of Example 1, except substituting the compounds of Preparation 7 or Preparation 3 for 3-(2-diisopropylamino)ethoxy-4-methoxyaniline and substituting 2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl]-4-carboxylic acid (EP 0533268) for 4-biphenylcarboxylic acid, gave the title compounds:

20 N-[3-(2-diisopropylamino)ethoxy-2-methylphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-[1,1'-biphenyl]-4-carboxamide: MS(ES) m/e 527.3 [M+H]⁺; and

N-[3-(2-diisopropylamino)ethoxy-phenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-[1,1'-biphenyl]-4-carboxamide: MS(ES) m/e 513.2 [M+H]⁺.

25

Example 14

Preparation of N-[3-(2-Dipropylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide,

30 Following the procedure of Example 1, except substituting the compound of Preparation 10(b) for 3-(2-diisopropylamino)ethoxy-4-methoxyaniline, gave the title compound MS(ES) m/e 447.1 [M+H]⁺.

Examples 15-22

35 Preparation of N-[3-(2-Dicyclohexylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide; N-[3-[2-(N-Cyclohexyl-N-isopropylamino)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-carboxamide; N-[3-[2-(cis-2,6-Dimethylpiperidin-1-yl)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide; N-[3-[2-(N-Ethyl-

- N-isopropylamino)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide; N-[3-[2-(2,5-Dimethylpyrrolidin-1-yl)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide; N-[4-Methoxy-3-[2-(2,2,6,6-tetramethylpiperidin-1-yl)ethoxy]phenyl]-1,1'-biphenyl-4-carboxamide; N-[3-[2-(N-tert-Butyl-N-isopropyl)aminoethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide; and N-[3-(2-Diisopropylamino)ethoxy-4-methylphenyl]-[1,1'-biphenyl]-4-carboxamide

- Following the procedure of Example 1, except substituting the compounds of Preparation 9, Preparations 11-16, or Preparation 8, for 3-(2-diisopropylamino)ethoxy-4-methoxyaniline, gave the title compounds:
- 10 N-[3-(2-dicyclohexylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide: MS(ES) m/e 527.5 [M+H]⁺;
- N-[3-[2-(N-cyclohexyl-N-isopropylamino)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-carboxamide: MS(ES) m/e 487.2 [M+H]⁺;
- N-[3-[2-(cis-2,6-dimethylpiperidin-1-yl)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide: MS(ES) m/e 459.0 [M+H]⁺;
- 15 N-[3-[2-(N-ethyl-N-isopropylamino)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide: MS(ES) m/e 433.1 [M+H]⁺;
- N-[3-[2-(2,5-dimethylpyrrolidin-1-yl)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide: MS(ES) m/e 445.0 [M+H]⁺;
- 20 N-[4-methoxy-3-[2-(2,2,6,6-tetramethylpiperidin-1-yl)ethoxy]phenyl]-1,1'-biphenyl-4-carboxamide: MS(ES) m/e 486.9 [M+H]⁺;
- N-[3-[2-(N-tert-butyl-N-isopropyl)aminoethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide: MS(ES) m/e 460.9 [M+H]⁺; and
- N-[3-(2-diisopropylamino)ethoxy-4-methylphenyl]-[1,1'-biphenyl]-4-carboxamide: MS(ES) m/e 431.0 [M+H]⁺.
- 25

Example 23

Preparation of N-[3-(3-Diisopropylamino)propyloxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide

- 30 Following the procedure of Example 1, except substituting the compound of Preparation 19(b) for 3-(2-diisopropylamino)ethoxy-4-methoxyaniline, gave the title compound: MS(ES) m/e 461.0 [M+H]⁺.

Example 24

- 35 Preparation of N-[3-(3-Diisopropylamino)propyl-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide Following the procedure of Example 1, except

substituting the compound of Preparation 20(c) for 3-(2-diisopropylamino)ethoxy-4-methoxyaniline, gave the title compound: MS(ES) m/e 445.2 [M+H]⁺.

Example 25

5 Preparation of N-[3-(2-Diisopropylamino)ethoxy-4-iodophenyl]-[1,1'-biphenyl]-4-carboxamide Following the procedure of Example 1, except substituting the compound of Preparation 21(b) for 3-(2-diisopropylamino)ethoxy-4-methoxyaniline, gave the title compound: MS(ES) m/e 542.9 [M+H]⁺.

Example 26

10 Preparation of N-[3-(2-Diisopropylamino)ethoxy-4-ethylphenyl]-[1,1'-biphenyl]-4-carboxamide

Following the procedure of Example 1, except substituting the compound of Preparation 22(c) for 3-(2-diisopropylamino)ethoxy-4-methoxyaniline, gave the
15 title compound: MS(ES) m/e 445.1 [M+H]⁺.

Example 27

Preparation of N-[3-(2-Diisopropylamino)ethoxy-4-(methoxycarbonylamino)phenyl]-[1,1'-biphenyl]-4-carboxamide

20 Following the procedure of Example 1, except substituting the compound of Preparation 23(c) for 3-(2-diisopropylamino)ethoxy-4-methoxyaniline, gave the title compound: MS(ES) m/e 490.1 [M+H]⁺.

Example 28

25 Preparation of N-[1,1'-Biphenyl]-4-yl-3-(2-diisopropylamino)ethoxy-4-methoxybenzamide a) N-[1,1'-biphenyl]-4-yl-3-hydroxy-4-methoxybenzamide

A solution of 3-hydroxy-4-methoxybenzoic acid (0.5 g, 3 mmol) in dichloromethane and dimethylformamide was cooled to -20°C, and N-bromosuccinimide (0.57 g, 3.2 mmol) and triphenylphosphine (0.8 g, 3.1 mmol)
30 were added in one portion. The mixture was stirred at -20°C for 10 min, and treated with a solution of 4-aminobiphenyl (0.5 g, 3 mmol) and dry pyridine (0.3 g, 3.8 mmol) in dichloromethane (5 mL). The mixture was allowed to warm to RT, stirred 20 h, concentrated *in vacuo*, and the residue was stirred in 5% methanol/dichloromethane (40 mL) and filtered. The filtrate was concentrated *in vacuo*, chromatographed (silica gel, 5% methanol/dichloromethane), and fractions
35 containing the product were pooled, concentrated *in vacuo*, and dissolved in

dichloromethane from which the title compound precipitated (0.11 g): MS(ES) m/e 320 [M+H]⁺.

b) N-[1,1'-biphenyl]-4-yl-3-(2-diisopropylamino)ethoxy-4-methoxybenzamide

- 5 Following the procedure of Preparation 1(a), except substituting the compound of Example 28(a) for 4-nitroguaiacole, gave the title compound: MS(ES) m/e 446.6 [M+H]⁺.

Example 29

- 10 Preparation of N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-methanamine Dihydrochloride

A solution of the compound of Example 1 (70 mg, 0.16 mmol) in anhydrous tetrahydrofuran (25 mL) was treated with 1.0M borane in tetrahydrofuran (1.56 mL, 1.56 mmol), heated to reflux for 3 h, cooled, maintained
15 at RT for 16 h, carefully treated with methanol (5 mL), stirred for 1 h, and concentrated *in vacuo*. The residue was partitioned between ethyl acetate (50 mL) and water (10 mL), and the organic phase was dried (Na₂SO₄), concentrated *in vacuo*, dissolved in dichloromethane, and treated with ethereal hydrogen chloride to afford the title compound(51 mg): MS(ES) m/e 433.3 [M+H]⁺.

20

Examples 30-31

- Preparation of N-[3-(2-Diisopropylamino)ethoxy-4-methoxy-phenyl]-4'-cyano-1,1'-biphenyl-4-carboxamide and N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4'-cyano-2'-methyl-1,1'-biphenyl-4-carboxamide Following the
25 procedure of Example 1, except substituting 4'-cyano-4-biphenylcarboxylic acid (WO 94/12181) or 4'-cyano-4'-methyl-4-biphenylcarboxylic acid (WO 96/19477) for 4-biphenylcarboxylic acid, gave the title compounds:

N-[3-(2-diisopropylamino)ethoxy-4-methoxy-phenyl]-4'-cyano-1,1'-biphenyl-4-carboxamide): MS(ES) m/e 472.0 [M+H]⁺; and

- 30 N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-4'-cyano-2'-methyl-1,1'-biphenyl-4-carboxamide): MS(ES) m/e 486.0 [M+H]⁺.

Example 32

- Preparation of N-[3-(2-Diisopropylamino)ethoxy-4-(ethoxycarbonyl)phenyl]-[1,1'-biphenyl]-4-carboxamide

35

Following the procedure of Example 1, except substituting 3-(2-diisopropylamino)ethoxy-4-(ethoxycarbonyl)aniline (Clinton, et al., *J. Am. Chem.*

Soc. 1957, 79, 2290) for 3-(2-diisopropylamino)ethoxy-4-methoxyaniline, gave the title compound: MS(ES) m/e 489.1 [M+H]⁺.

Example 33

- 5 Preparation of N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-2-naphthalenecarboxamide Following the procedure of Example 1, except substituting 2-naphthalenecarboxylic acid for 4-biphenylcarboxylic acid, gave the title compound: MS(ES) m/e 421.3 [M+H]⁺.

Example 34

- 10 Preparation of N-(1'-Methylspiro[benzofuran-3(2H),4'-piperidin]-5-yl-[1,1'-biphenyl]-4-carboxamide Following the procedure of Example 1, except substituting 1'-methyl-spiro[benzofuran-3(2H),4'-piperidin]-5-amine (WO 96/11934) for 3-(2-diisopropylamino)ethoxy-4-methoxyaniline, gave the title
15 compound: MS(ES) m/e 399.2 [M+H]⁺.

Examples 35-36

- Preparation of N-[3-[2-(N-Isopropyl-N-methylamino)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide and N-Ethyl-N-[3-[2-(N-isopropyl-N-methylamino)ethoxy]-4-methoxyphenyl]-1,1'-biphenyl-4-carboxamide
20

Following the procedure of Example 1, except substituting the compounds of Preparations 17(b) and 18(b) for 3-(2-diisopropylamino)ethoxy-4-methoxyaniline gave the title compounds:

- 25 N-[3-[2-(N-isopropyl-N-methylamino)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide: MS(ES) m/e 419.1 [M+H]⁺; and
N-ethyl-N-[3-[2-(N-isopropyl-N-methylamino)ethoxy]-4-methoxyphenyl]-1,1'-biphenyl-4-carboxamide: MS(ES) m/e 446.3 [M+H]⁺.

Example 37

- 30 Preparation of N-Methyl-N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-[1,1'-biphenyl]-4-carboxamide

- Following the procedure of Example 1, except substituting the compound of Preparation 24(b) for 3-(2-diisopropylamino)ethoxy-4-methoxyaniline and substituting 2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carboxylic
35 acid (EP 0533268) for 4-biphenylcarboxylic acid, gave the title compound: MS(ES) m/e 557.3 [M+H]⁺.

Example 38

Preparation of N-Methyl-N-[3-(diisopropylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide

Following the procedure of Example 1, except substituting the compound of Preparation 24(b) for 3-(2-diisopropylamino)ethoxy-4-methoxyaniline, gave the title compound: MS(ES) m/e 461.3 [M+H]⁺.

Example 39

Preparation of N-[4-(2-Diisopropylamino)ethoxy-3-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide

a) resin-bound 4'-(4-biphenylcarboxylic acid)

Resin-bound phenylbromide (WO 98/17695, 3-Scheme18)(1.0 g, 1.22 mmol) was swelled in a mixture of toluene (10 mL) and ethanol (4 mL) for 30 min, and treated with 4-carboxybenzeneboronic acid (0.6 g, 3.6 mmol), tetrakis-(triphenylphosphine)palladium(0) (0.2 g), and 2M aqueous sodium carbonate (3.6 mL, 7.2 mmol). The mixture heated to reflux for 24 h, cooled, and the resin was filtered and washed sequentially with dichloromethane, methanol, 1:1 methanol/water containing 15 drops of concentrated hydrochloric acid, 1:1 methanol/water, dichloromethane, and methanol. The resulting resin was dried *in vacuo*.

b) resin-bound 4'-(biphenyl-4-carbonyl chloride)

The resin of Example 39(a) (0.06 g) and oxalyl chloride (0.13 g, 1 mmol) in toluene (10 mL) was shaken for 16 h, filtered, and washed with toluene and dichloromethane.

c) resin bound 4'-[N-[4-(2-diisopropylamino)ethoxy-3-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide]

A mixture of the resin of Example 39(b), the compound of Preparation 1(b)(0.13 g, 0.5 mmol), and diisopropylethylamine (0.05 g, 0.4 mmol) in dichloromethane (15 mL) was shaken for 20 h, the resin was filtered, washed with dichloromethane (3 x 20 mL) and with methanol (2 x 20 mL), and dried *in vacuo* for 16 h.

d) N-[4-(2-diisopropylamino)ethoxy-3-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide

A mixture of the resin of Example 39(c), trifluoroacetic acid (3 mL) and dichloromethane (0.5 mL) was stirred for 20 h, and the resin was filtered and washed with methanol (3 x 10 mL). The filtrate was concentrated *in vacuo*, and the residue was partitioned between dichloromethane (50 mL) and 10% aqueous

sodium hydroxide (5 mL). The organic phase was washed with water, dried (Na_2SO_4), and concentrated *in vacuo* to give the title compound: MS(ES) m/e 447.4 $[\text{M}+\text{H}]^+$.

5

Examples 40-44

Preparation of N-[2-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide; N-[3-(2-Diisopropylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide; N-[2-Fluoro-4-(2-diisopropylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide; N-[4-(2-Diisopropylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide; and N-[2-(2-Diisopropylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide

10

Following the procedure of Example 39(c)-39(d), except substituting the compounds of Preparations 2-6 for the compound of Preparation 1, gave the title compounds:

15

N-[2-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide: MS(ES) m/e 447.3 $[\text{M}+\text{H}]^+$;

N-[3-(2-diisopropylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide: MS(ES) m/e 417.5 $[\text{M}+\text{H}]^+$;

20

N-[2-fluoro-4-(2-diisopropylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide: MS(ES) m/e 435.4 $[\text{M}+\text{H}]^+$;

N-[4-(2-diisopropylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide MS(ES) m/e 417.4 $[\text{M}+\text{H}]^+$; and

N-[2-(2-diisopropylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide: MS(ES) m/e 417.4 $[\text{M}+\text{H}]^+$.

25

Example 45

Preparation of N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4-(iodo)benzamide

a) [3-(2-diisopropylamino)ethoxy-4-methoxyaniline/(4-formyl-3,5-dimethoxyphenoxy)-Merrifield resin adduct

30

A mixture of 4-formyl-3,5-dimethoxyphenoxy-Merrifield resin (Booamra et al., *J. Org. Chem.* 1995, 60, 5742), 3-(2-diisopropylamino)ethoxy-4-methoxyaniline (WO 9515954), and sodium triacetoxymethylborohydride in dimethylformamide containing 1% acetic acid was shaken to afford the title adduct.

35

b) N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-4-iodobenzamide/(4-formyl-3,5-dimethoxyphenoxy)-Merrifield resin adduct

The resin of Example 45(a) in dimethylformamide was treated with pyridine and with an excess of a mixture of equivalent amounts of 4-iodobenzoic acid, N-bromosuccinimide and triphenylphosphine. The resin was washed to afford the title adduct.

5 c) N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-4-iodobenzamide

The resin of Example 45(b) was stirred in a mixture of trifluoroacetic acid:dichloromethane:water (50:48:2), filtered, and the filtrate concentrated *in vacuo* to afford the title compound: MS(ES) m/e 497.8 [M+H]⁺.

10

Examples 46-70

Following the procedure of Example 45, except using 3-(2-diethylamino)ethoxy-aniline (WO 95/15954), 4-(2-diethylamino)ethoxy-aniline (Wyatt et al., *J. Med. Chem.*, 1995, 38, 1657), 3-(2-diethylamino)ethoxy-4-methoxyaniline (WO 96/23769), 3-(2-diisopropylamino)ethoxy-4-methoxyaniline (WO 95/15954), and the compounds of Preparation 3 and Preparation 5, and using 4-isopropylbenzoic acid, 4-cyclohexylbenzoic acid, 4'-ethyl-4-biphenylcarboxylic, 4-nitrobenzoic acid, 4-methoxybenzoic acid, 3-bromobenzoic acid, 3-iodobenzoic acid, 4-bromobenzoic acid, and 4-iodobenzoic acid gave the following compounds:

20 N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-4-iodobenzamide: MS(ES) m/e 497.0 [M+H]⁺;

N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-4-bromobenzamide: MS(ES) m/e 449.0 [M+H]⁺;

N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-3-bromobenzamide: MS(ES) m/e 449.0 [M+H]⁺;

25 N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-4-(isopropyl)benzamide: MS(ES) m/e 413.2 [M+H]⁺;

N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-4'-ethyl-[1,1'-biphenyl]-4-carboxamide: MS(ES) m/e 475.2 [M+H]⁺;

30 N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-4-(methoxy)benzamide: MS(ES) m/e 401.3 [[M+H]⁺];

N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-4-(nitro)benzamide: MS(ES) m/e 416.5 [M+H]⁺;

N-[4-(2-diisopropylamino)ethoxy-phenyl]-4-iodobenzamide: MS(ES) m/e 467.0 [M+H]⁺;

35 N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-3-iodobenzamide: MS(ES) m/e 497.0 [M+H]⁺;

- N-[3-(2-diethylamino)ethoxy-4-methoxyphenyl]-3-iodobenzamide:
MS(ES) m/e 460.0 [M+H]⁺;
- N-[4-(2-diisopropylamino)ethoxy-phenyl]-3-iodobenzamide: MS(ES) m/e
467.0 [M+H]⁺;
- 5 N-[3-(2-diisopropylamino)ethoxy-phenyl]-3-iodobenzamide: MS(ES) m/e
476.0 [M+H]⁺;
- N-[3-(2-diethylamino)ethoxy-phenyl]-3-iodobenzamide: MS(ES) m/e 439.0
[M+H]⁺;
- N-[3-(2-diethylamino)ethoxy-4-methoxyphenyl]-4-bromobenzamide:
10 MS(ES) m/e 423.0 [M+H]⁺;
- N-[4-(2-diisopropylamino)ethoxy-phenyl]-4-bromobenzamide: MS(ES)
m/e 418.8 [M+H]⁺;
- N-[3-(2-diethylamino)ethoxy-4-methoxyphenyl]-3-bromobenzamide:
MS(ES) m/e 423.0 [M+H]⁺;
- 15 N-[4-(2-diisopropylamino)ethoxy-phenyl]-3-bromobenzamide: MS(ES)
m/e 418.8 [M+H]⁺;
- N-[4-(2-diisopropylamino)ethoxy-phenyl]-4-(isopropyl)benzamide:
MS(ES) m/e 383.2 [M+H]⁺;
- N-[3-(2-diethylamino)ethoxy-4-methoxyphenyl]-4-(cyclohexyl)benzamide:
20 MS(ES) m/e 425.2 [M+H]⁺;
- N-[4-(2-diisopropylamino)ethoxy-phenyl]-4-(cyclohexyl)benzamide:
MS(ES) m/e 423.0 [M+H]⁺;
- N-[4-(2-diethylamino)ethoxy-phenyl]-4-(cyclohexyl)benzamide: MS(ES)
m/e 395.0 [M+H]⁺;
- 25 N-[3-(2-diethylamino)ethoxy-4-methoxyphenyl]-4'-ethyl-[1,1'-biphenyl]-4-
carboxamide: MS(ES) m/e 447.2 [M+H]⁺;
- N-[4-(2-diisopropylamino)ethoxy-phenyl]-4'-ethyl-[1,1'-biphenyl]-4-
carboxamide: MS(ES) m/e 445.2 [M+H]⁺;
- N-[4-(2-diethylamino)ethoxy-phenyl]-4'-ethyl-[1,1'-biphenyl]-4-
30 carboxamide: MS(ES) m/e 417.4 [M+H]⁺; and
- N-[3-(2-diisopropylamino)ethoxy-phenyl]-4'-ethyl-[1,1'-biphenyl]-4-
carboxamide: MS(ES) m/e 445.2 [M+H]⁺.

Examples 71-73

- 35 Preparation of N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-3,4-
dichlorobenzamide, N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-3,5-

dichlororbenzamide, and N-[3-(2-Diisopropylamino)ethoxy-4-methoxy-phenyl]-4-(cyclohexyl)benzamide

Following the procedure of Example 1, except substituting 3,4-dichlorobenzoic acid, 3,5-dichlorobenzoic acid or 4-cyclohexylbenzoic acid for 4-biphenylcarboxylic acid, gave the title compounds:

N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-3,4-dichlororbenzamide: MS(ES) m/e 439.3 [M+H]⁺;

N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-3,5-dichlororbenzamide: MS(ES) m/e 439.2 [M+H]⁺; and

10 N-[3-(2-diisopropylamino)ethoxy-4-methoxy-phenyl]-4-(cyclohexyl)benzamide: MS(ES) m/e 453.0 [M+H]⁺.

Example 74

Preparation of N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-3'-methoxy-[1,1'-biphenyl]-4-carboxamide

15 a) N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzamide/(4-formyl-3,5-dimethoxyphenoxy)-Merrifield resin adduct

Following the procedure of Example 45(b), except substituting 4-(4,4,5,5-tetramethyl-1,3,2-dioxaborolan-2-yl)benzoic acid (Guiles et al., *J. Org. Chem.* 1996, 61, 5169) for 4-iodobenzoic acid, gave the title adduct.

20 b) N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-3'-methoxy-[1,1'-biphenyl]-4-carboxamide/(4-formyl-3,5-dimethoxyphenoxy)-Merrifield resin adduct

Following the general procedure of Guiles et al., *J. Org. Chem.* 1996, 61, 5169, the adduct of Example 74(a) in dimethylformamide was reacted with 3-iodoanisole, tetrakis(triphenylphosphine)palladium(0), and aqueous potassium carbonate at 80°C for 18 h to afford the title adduct.

c) N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-3'-methoxy-[1,1'-biphenyl]-4-carboxamide

30 Following the procedure of Example 45(c), except substituting the adduct of Example 74(b) for the adduct of Example 42(b), gave the title compound: MS(ES) m/e 477.2 [M+H]⁺.

Examples 75-81

35 Preparation of N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-4'-methoxy-[1,1'-biphenyl]-4-carboxamide; N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-4'-hydroxy-[1,1'-biphenyl]-4-carboxamide; N-[3-(2-

Diisopropylamino)ethoxy-4-methoxyphenyl]-4'-acetyl-[1,1'-biphenyl]-4-carboxamide N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-3'-methoxy-[1,1'-biphenyl]-4-carboxamide N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-3'-methoxy-[1,1'-biphenyl]-4-carboxamide N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-3'-hydroxy-[1,1'-biphenyl]-4-carboxamide and N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-3'-hydroxy-[1,1'-biphenyl]-4-carboxamide

Following the procedure of Example 74, except using 3-(2-diisopropylamino)ethoxy-4-methoxyaniline (WO 95/15954), 3-(2-diethylamino)ethoxy-4-methoxyaniline, and 3-(2-dimethylamino)ethoxy-4-methoxyaniline, and using 3-iodophenol, 3-iodoanisole, and 4'-iodoacetophenone, gave the title compounds:

N-[3-(2-dimethylamino)ethoxy-4-methoxyphenyl]-4'-methoxy-[1,1'-biphenyl]-4-carboxamide MS(ES) m/e 421.0 [M+H]⁺;
 N-[3-(2-dimethylamino)ethoxy-4-methoxyphenyl]-4'-hydroxy-[1,1'-biphenyl]-4-carboxamide MS(ES) m/e 407.2 [M+H]⁺;
 N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-4'-acetyl-[1,1'-biphenyl]-4-carboxamide: MS(ES) m/e 489.2 [M+H]⁺;
 N-[3-(2-diethylamino)ethoxy-4-methoxyphenyl]-3'-methoxy-[1,1'-biphenyl]-4-carboxamide: MS(ES) m/e 449.2 [M+H]⁺;
 N-[3-(2-dimethylamino)ethoxy-4-methoxyphenyl]-3'-methoxy-[1,1'-biphenyl]-4-carboxamide: MS(ES) m/e 421.0 [M+H]⁺;
 N-[3-(2-diisopropylamino)ethoxy-4-methoxyphenyl]-3'-hydroxy-[1,1'-biphenyl]-4-carboxamide: MS(ES) m/e 463.2 [M+H]⁺; and
 N-[3-(2-dimethylamino)ethoxy-4-methoxyphenyl]-3'-hydroxy-[1,1'-biphenyl]-4-carboxamide MS(ES) m/e 407.2 [M+H]⁺.

Example 82

Preparation of N-[2-(Dimethylaminomethyl)-1,4-benzodioxan-7-yl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carboxamide

The compound of Preparation 26(c) (188 mg, 0.6 mmol) in tetrahydrofuran (30 mL) was treated with a solution of sodium hydroxide (55 mg) in water (1 mL) and 2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-biphenyl-4-carbonyl chloride (WO 95/15954) (0.6 mmol) was added. The mixture was stirred for 16 h, concentrated *in vacuo*, and partitioned between water and chloroform. The organic phase was dried (Na₂SO₄) and concentrated *in vacuo* to give a white foam which

was chromatographed (basic alumina, ethyl acetate) to give the title compound.
mp 92-94°C.

Example 84

5 Preparation of N-[2-(dimethylaminomethyl)-1,4-benzodioxan-6-yl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-

Following the procedure of Example 82, except substituting the compound of Preparation 25(c) for the compound of Preparation 26(c), gave the title compound. mp 188-190°C.

10

Biological Data:

CCR5 Receptor Binding Assay

CHO cell membranes (0.25 x 10⁶ cell equivalents) derived from CHO cells stably transfected with CCR5 were incubated with 0.3 ¹²⁵I-RANTES in a 96 well plate
15 for 45 min. at room temperature (final reaction volume 200 ul). The reaction was terminated by filtration and the filters (GF/C) were washed twelve times with a solution of phosphate buffered saline containing 0.1 % bovine serum albumin and 0.05 % NaN₃. The radioactivity bound to filters was measured by liquid
20 scintillation spectrometry. Non-specific binding was determined in the presence of unlabelled RANTES (10 or 30 nM) and averages 30-50% of total binding.

CCR5 Receptor Functional Assay

The cellular functional assay used to assess antagonist activity of compounds was RANTES-induced Ca²⁺ mobilization in RBL 2H3 cells stably expressing the
25 hCCR5 receptor (RBL 2H3 hCCR5). Agonist activity is determined by Ca²⁺ mobilization in the same cells which is inhibitable by a selective CCR5 antagonist. Cells were grown to 80-100% confluency in T-150 flasks and washed with phosphate-buffered saline. Cells were lifted from the flasks by treating with 3 mL of 1 mM EDTA for 3 min. at room temperature and diluting to 2 X 10⁶ cells/mL
30 with Krebs Ringer Henseleit buffer (KRH; 118 mM NaCl, 4.6 mM KCl, 25 mM NaHCO₃, 1 mM KH₂PO₄ and 11 mM glucose) containing 5 mM HEPES (pH 7.4), 1 mM CaCl₂, 1 mM MgCl₂ and 0.1% BSA and centrifuged at 200g for 3 min. Cells were resuspended at 2 X 10⁶ cells/mL in the same buffer with 2 µM Fura-2AM, and incubated for 35 min. at 37° C. Cells were centrifuged at 200 x g
35 for 3 min. and resuspended in the same buffer without Fura-2AM, then incubated for 15 min. at 37° C to complete the hydrolysis of intracellular Fura-2AM, and then centrifuged as before. Cells (10⁶ cells/mL) were resuspended in cold KRH

with 5 mM HEPES (pH 7.4), 1 mM CaCl_2 , 1 mM MgCl_2 and 0.1% gelatin and maintained on ice until assayed. For antagonist studies, aliquots (2 mL) of cells were prewarmed at 37° C for 5 min. in 3 mL plastic cuvettes and fluorescence measured in a fluorometer (Johnson Foundation Biomedical Group, Philadelphia, PA, USA) with magnetic stirring and temperature maintained at 37° C. Excitation was set at 340 nm and emission set at 510 nm. Various concentrations of antagonists or vehicle were added and fluorescence monitored for ~15 sec to ensure that there was no change in baseline fluorescence, followed by the addition of 33 nM RANTES. Maximal Ca^{2+} attained after 33 nM RANTES stimulation was calculated as described by Grynkiewicz *et al.*, (1985). The percent of maximal RANTES-induced Ca^{2+} was determined for each concentration of antagonist and the IC_{50} , defined as the concentration of test compound that inhibits 50% of the maximal 33 nM RANTES response, obtained from the concentration-response curves (5-7 concentrations of antagonists).

The compounds of this invention show CCR5 receptor ligand activity having IC_{50} values in the range of 0.0001 to 100 μM . The full structure/activity relationship has not yet been established for the compounds of this invention. However, given the disclosure herein, one of ordinary skill in the art can utilize the present assays in order to determine which compounds of formula (I) are ligands of the CCR5 receptor and which bind thereto with an IC_{50} value in the range of 0.0001 to 100 μM .

All publications, including, but not limited to, patents and patent applications cited in this specification, are herein incorporated by reference as if each individual publication were specifically and individually indicated to be incorporated by reference herein as though fully set forth.

The above description fully discloses the invention including preferred embodiments thereof. Modifications and improvements of the embodiments specifically disclosed herein are within the scope of the following claims. Without further elaboration it is believed that one skilled in the art can, given the preceding description, utilize the present invention to its fullest extent. Therefore any examples are to be construed as merely illustrative and not a limitation on the scope of the present invention in any way. The embodiments of the invention in which an exclusive property or privilege is claimed are defined as follows.


What is claimed is:

1. A method of treating a CCR5-mediated disease state in mammals which comprises administering to a mammal in need of such treatment, an effective amount of a compound of formula (I) or a pharmaceutically acceptable salt thereof:



Formula I

in which Ar represents a group selected from (i), (ii) or (iii);



in which:

R^1 and R^2 are independently hydrogen, C_{1-6} alkyl, C_{2-6} alkenyl, C_{2-6} alkynyl, C_{3-7} cycloalkyl, C_{3-6} cycloalkenyl, aryl, $(CH_2)_aNR^7R^8$, $(CH_2)_aNR^7COR^9$, $(CH_2)_aNR^7CO_2R^{10}$, $(CH_2)_aNR^7SO_2R^{11}$, $(CH_2)_aCONR^{12}R^{13}$, hydroxy C_{1-6} alkyl, C_{1-4} alkoxyalkyl (optionally substituted by a C_{1-4} alkoxy or hydroxy group), $(CH_2)_aCO_2C_{1-6}$ alkyl, $(CH_2)_bOC(O)R^{14}$, $CR^{15}=NOR^{16}$, $CNR^{15}=NOR^{16}$, COR^{17} , $CONR^{12}R^{13}$, $CONR^{12}(CH_2)_cOC_{1-4}$ alkyl, $CONR^{12}(CH_2)_aCO_2R^{18}$, $CONHNR^{19}R^{20}$, $CONR^{12}SO_2R^{21}$, CO_2R^{22} , cyano, trifluoromethyl, NR^7R^8 , NR^7COR^9 , $NR^{23}CO(CH_2)_aNR^{23}R^{24}$, $NR^{23}CONR^{23}R^{24}$, $NR^7CO_2R^{10}$, $NR^7SO_2R^{11}$, $N=CNR^{23}NR^{23}R^{24}$, nitro, hydroxy, C_{1-6} alkoxy, hydroxy C_{1-6} alkoxy, C_{1-6} alkoxy C_{1-6} alkoxy, $OC(O)NR^{25}R^{26}$, SR^{27} , SOR^{28} , SO_2R^{28} , $SO_2NR^{25}R^{26}$ or halogen;

a is 1, 2, 3 or 4;

R^7 and R^8 are independently hydrogen or C_{1-6} alkyl, or NR^7R^8 forms a heterocyclic ring which has 5 or 6 ring members which, may optionally be substituted by an oxo group and, when there are 6 ring members, may optionally contain in the ring one oxygen or sulfur atom;

R^9 is hydrogen, C_{1-6} alkyl or C_{1-4} alkoxyalkyl;

R^{10} is C_{1-6} alkyl;

R^{11} is C_{1-6} alkyl or phenyl;

R¹² and R¹³ are independently hydrogen or C₁₋₆alkyl, or NR¹²R¹³ forms a saturated heterocyclic ring which has 5 or 6 members which, when there are 6 ring members, may optionally contain in the ring one oxygen or sulfur atom;

b is 0, 1, 2 or 3;

R¹⁴ is C₁₋₄alkyl, optionally substituted by a C₁₋₆alkoxy;

R¹⁵ and R¹⁶ are independently hydrogen or C₁₋₆alkyl;

R¹⁷ is hydrogen or C₁₋₆alkyl;

c is 1, 2 or 3;

R¹⁸ is hydrogen or C₁₋₆alkyl;

R¹⁹ and R²⁰ are independently hydrogen or C₁₋₆alkyl;

R²¹ is hydrogen or C₁₋₆alkyl;

R²² is hydrogen or C₁₋₆alkyl optionally substituted with one or two substituents selected from C₁₋₆alkyl, C₁₋₆alkoxy, hydroxy, or NR⁷R⁸;

R²³ and R²⁴ are independently hydrogen or C₁₋₆alkyl;

R²⁵ and R²⁶ are independently hydrogen or C₁₋₆alkyl, or NR²⁵R²⁶ forms a saturated heterocyclic ring which has 5 or 6 members which, when there are 6 ring members, may optionally contain in the ring one oxygen or sulfur atom;

R²⁷ is hydrogen or C₁₋₆alkyl;

R²⁸ is C₁₋₆alkyl;

P is a 5 to 7-membered heterocyclic ring containing 1 to 4 heteroatoms selected from oxygen, nitrogen or sulfur;

R³ and R⁴ are independently hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl, C₃₋₆cycloalkenyl, hydroxyC₁₋₆alkyl, C₁₋₆alkylOC₁₋₆alkyl, CONR²⁹R³⁰, CO₂R³¹, cyano, aryl, trifluoromethyl, NR²⁹R³⁰, nitro, hydroxy, C₁₋₆alkoxy, acyloxy or halogen;

R²⁹, R³⁰ and R³¹ are independently hydrogen or C₁₋₆alkyl;

R⁵ is hydrogen, C₁₋₆alkyl, C₁₋₆alkoxy or halogen;

R⁶ is hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl (optionally substituted by a hydroxy or an oxo group), hydroxyC₁₋₆alkyl, hydroxyC₃₋₆alkenyl, hydroxyC₃₋₆alkynyl, (CH₂)_dOR³², (CH₂)_dCOR³³, (CH₂)_dCR³⁴=NOR³⁵, CONR³⁶R³⁷, CO₂R³⁸, hydroxy, O(CH₂)_eR³⁹, NR³⁶R³⁷, SR⁴⁰, SO₂NR⁴¹R⁴² or halogen;

d is 0, 1, 2, 3, 4, 5, or 6;

R³² is C₁₋₆alkyl, hydroxyC₁₋₆alkyl, or C₁₋₄alkanoyl;

R³³ is hydrogen or C₁₋₆alkyl;

R³⁴ is hydrogen or C₁₋₆alkyl;

R³⁵ is hydrogen or C₁₋₆alkyl;

R^{36} and R^{37} are independently hydrogen or C_{1-6} alkyl or $NR^{36}R^{37}$ forms a saturated heterocyclic ring which has 5 or 6 members, may optionally be substituted by an oxo group and, when there are 6 ring members, may optionally contain one oxygen or sulfur atom or an NH or NR^{43} where R^{43} is C_{1-6} alkyl, COR^{44} or CO_2R^{45} where R^{44} and R^{45} are independently hydrogen or C_{1-6} alkyl;

R^{38} is hydrogen or C_{1-6} alkyl;

e is 1, 2, 3, 4, 5 or 6;


R^{39} is C_{1-6} alkoxy, CO_2H , CO_2C_{1-6} alkyl or $CONR^{36}R^{37}$;

R^{40} is C_{1-6} alkyl;

R^{41} and R^{42} are independently hydrogen or C_{1-6} alkyl;

alternatively, R^5 and R^6 form a fused benzo ring optionally substituted with C_{1-6} alkyl, C_{1-6} alkoxy or halogen;

when Ar is (i), (ii) or (iii), and A is $CONR^{46}$, $NHCO$, $-NHCH_2$, or CH_2NH , where R^{46} is hydrogen or C_{1-6} alkyl, E represents (a):



(a);

in which

R^{47} and R^{48} are independently hydrogen or C_{1-6} alkyl;

R^{49} is hydrogen, C_{1-6} alkyl, CO_2R^{53} , $NHCO_2R^{54}$, hydroxy, C_{1-6} alkoxy or halogen where R^{53} is hydrogen or C_{1-6} alkyl and R^{54} is C_{1-6} alkyl;

R^{50} and R^{51} are independently hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl, aralkyl, or together with the nitrogen atom to which they are attached form an optionally substituted 5- to 7-membered heterocyclic ring containing one to two heteroatoms selected from oxygen, nitrogen or sulfur;


B is oxygen, $S(O)_h$ where h is 0, 1 or 2, $CR^{55}=CR^{56}$ or $CR^{55}R^{56}$ where R^{55} and R^{56} are independently hydrogen or C_{1-6} alkyl, or B is NR^{57} where R^{57} is hydrogen, C_{1-6} alkyl or phenyl C_{1-6} alkyl;

R^{52} is hydrogen or R^{52} taken together with R^{46} forms a group D where D is $(CR^{58}R^{59})_i$ where i is 2, 3 or 4 and R^{58} and R^{59} are independently hydrogen or C_{1-6} alkyl or D is $(CR^{58}R^{59})_j-G$ where j is 0, 1, 2 or 3 and G is oxygen, sulfur or $CR^{58}=CR^{59}$;

f is 1 to 4; and

g is 1 or 2;

when Ar is (i), (ii) or (iii), and A is $CONR^{60}$, $NHCO$, or CH_2NH , where R^{60} is hydrogen or C_{1-6} alkyl, E represents (b):



(b);

R^{61} is hydrogen or C_{1-6} alkyl or R^{61} and R^{60} together form a group -K- where K is $(CR^{65}R^{66})_m$ where m is 2, 3, or 4 and R^{65} and R^{66} are independently hydrogen or C_{1-6} alkyl or K is $(CR^{65}R^{66})_n-L$ where n is 0, 1, 2, or 3 and L is oxygen, sulfur or $CR^{65}=CR^{66}$;

R^{62} is hydrogen or C_{1-6} alkyl;


R^{63} and R^{64} are independently hydrogen or C_{1-6} alkyl;

J is oxygen, $CR^{67}R^{68}$ or NR^{69} where R^{67} , R^{68} and R^{69} are independently hydrogen or C_{1-6} alkyl or J is a group $S(O)_m$ where m is 0, 1 or 2;

k is 1, 2 or 3;

l is 1, 2 or 3;

when Ar is (i), (ii) or (iii), and A is $CONR^{70}$, $NHCO$, $-NHCH_2$, or CH_2NH , where R^{70} is hydrogen or C_{1-6} alkyl, E represents (c):



(c);

in which:

M is oxygen, $S(O)_p$ where p is 0, 1 or 2, $CR^{76}=CR^{77}$ or $CR^{76}R^{77}$ where R^{76} and R^{77} are independently hydrogen or C_{1-6} alkyl, or M is NR^{78} where R^{78} is hydrogen or alkyl;

R^{71} and R^{72} are independently hydrogen or C_{1-6} alkyl;


R^{73} is hydrogen, C_{1-6} alkyl, CO_2R^{79} , $NHCO_2R^{80}$, hydroxy, C_{1-6} alkoxy or halogen where R^{79} is hydrogen or C_{1-6} alkyl and R^{80} is C_{1-6} alkyl;

R^{74} is hydrogen or together with R^{70} forms a group Q where Q is $CR^{81}=CR^{82}$, $CR^{81}=CR^{82}CR^{81}R^{82}$ or $(CR^{81}R^{82})_q$ where q is 2 or 3 and R^{81} and R^{82} are independently hydrogen or C_{1-6} alkyl;


n is 0, 1, 2 or 3;

o is 1 or 2;

R^{75} is a group of formula (d):




where r, s and t are independently integers having the value 1, 2 or 3;
or R⁷⁵ is a group of formula (e):



where u is 0, 1, 2 or 3 and R⁸³ is hydrogen or C₁₋₆alkyl;

when Ar is (i), (ii), or (iii), and A is CONH, NHCO or CH₂NH, E represents (f):




R⁸⁴ and R⁸⁵ are independently hydrogen or C₁₋₆alkyl;

R⁸⁶ and R⁸⁷ are independently hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl, aralkyl, or together with the nitrogen atom to which they are attached form an optionally substituted 5- to 7-membered heterocyclic ring containing one to two heteroatoms selected from oxygen, nitrogen or sulfur;

T is -(CR⁸⁸R⁸⁹)_w- or -O(CR⁸⁸R⁸⁹)_x- where R⁸⁸ and R⁸⁹ are independently hydrogen or C₁₋₆alkyl where w is 2 or 3 and x is 1, 2 or 3;
v is 1 to 4; and

W is oxygen, S(O)_y where y is 0, 1 or 2, or W is NR⁹⁰ where R⁹⁰ is hydrogen or C₁₋₆alkyl, or W is CR⁹¹=CR⁹² or CR⁹¹R⁹² where R⁹¹ and R⁹² are independently hydrogen or C₁₋₆alkyl;

when Ar is (i), (ii) or (iii), and A is CONR⁹³, NHCO, or CH₂NH where R⁹³ is hydrogen or C₁₋₆alkyl, E represents a group (g):



R⁹⁴ is hydrogen, halogen, hydroxy, C₁₋₆alkyl or C₁₋₆alkoxy or R⁹⁴ and R⁹³ taken together from a group -X- where X is (CR⁹⁷R⁹⁸)_{aa} where aa is 2, 3 or 4


and R^{97} and R^{98} are independently hydrogen or C_{1-6} alkyl or X is $(CR^{97}R^{98})_{ab}$ -Y where ab is 0, 1, 2 or 3 and Y is oxygen, sulfur or $CR^{97}=CR^{98}$;

R^{95} is hydrogen, C_{1-6} alkyl, CO_2R^{99} , $NHCO_2R^{100}$, hydroxy, C_{1-6} alkoxy or halogen where R^{95} is hydrogen or C_{1-6} alkyl and R^{100} is C_{1-6} alkyl;

z is 1 or 2; and

R^{96} is an optionally substituted 5 to 7-membered saturated or partially saturated heterocyclic ring containing 1 to 3 heteroatoms selected from nitrogen, oxygen or sulfur or R^{96} is an optionally substituted 6,6 or 6,5 bicyclic ring containing a nitrogen atom and optionally a further heteroatom selected from oxygen, nitrogen or sulfur;

when Ar is (i), (ii) or (iii) and A is $CONR^{101}$, $NHCO$, or CH_2NH , where R^{101} is hydrogen or C_{1-6} alkyl, E represents group (h):



R^{102} is hydrogen or C_{1-6} alkyl, C_{1-6} alkoxy or halogen, or R^{102} together with R^{101} form a group -AA- where AA is $(CR^{107}R^{108})_{ad}$ where ad is 1, 2 or 3 and R^{107} and R^{108} are independently hydrogen or C_{1-6} alkyl or AA is $(CR^{107}=CR^{108})_{ae}$ -AB where ae is 0, 1 or 2 and AB is oxygen, sulfur, $CR^{107}=CR^{108}$, $CR^{107}=N$, $CR^{102}NR^{108}$ or $N=N$;


R^{103} and R^{104} are independently hydrogen or C_{1-6} alkyl;

R^{105} and R^{106} are independently hydrogen, C_{1-6} alkyl, C_{3-7} cycloalkyl, aralkyl, or together with the nitrogen atom to which they are attached form an optionally substituted 5- to 7-membered heterocyclic ring containing one to two heteroatoms selected from oxygen, nitrogen or sulfur;

ac is 0 to 4;

Z is an optionally substituted 5 to 7-membered heterocyclic ring containing 1 to 3 heteroatoms selected from oxygen, nitrogen or sulfur;

when Ar is (i), (ii) or (iii) and A is $CONR^{109}$, $NHCO$, or CH_2NH where R^{109} is hydrogen or C_{1-6} alkyl, E represents group (i):



R¹¹⁰ is hydrogen or C₁₋₆alkyl or R¹¹⁰ and R¹⁰⁹ together form a group -AD- where AD is (CR¹¹⁵R¹¹⁶)_{ah} where ah is 2, 3 or 4 and R¹¹⁵ and R¹¹⁶ are independently hydrogen or C₁₋₆alkyl or AD is (CR¹¹⁵R¹¹⁶)_{ai}-AE where ai is 0, 1, 2 or 3 and AE is oxygen, sulfur or CR¹¹⁵=CR¹¹⁶;

R¹¹¹ and R¹¹² are independently hydrogen, C₁₋₆alkyl, C₃₋₇cycloalkyl, aralkyl, or together with the nitrogen atom to which they are attached form an optionally substituted 5- to 7-membered heterocyclic ring containing one to two heteroatoms selected from oxygen, nitrogen or sulfur;

R¹¹³ and R¹¹⁴ are independently hydrogen or C₁₋₆alkyl;

AC is oxygen, CR¹¹⁷R¹¹⁸ or NR¹¹⁹ where R¹¹⁷, R¹¹⁸ and R¹¹⁹ are independently hydrogen or C₁₋₆alkyl or AC is a group S(O)_{aj} where aj is 0, 1 or 2; af is 1, 2 or 3;

ag is 1, 2, 3, or 4; and

ah is 0, 1, 2, 3 or 4.

2. The method as claimed in claim 1 wherein the compound of formula (I) is a compound selected from:

N-[3-(3-Dimethylamino)propoxy-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-(1,1'-biphenyl)-4-carboxamide;

N-[3-(2-Piperidine)ethoxy-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-(1,1'-biphenyl)-4-carboxamide;

N-[3-(3-Dimethylaminopropyl)-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carboxamide oxalate;

N-[3-(1-Methyl-4-piperidyl)-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carboxamide oxalate;

N-[3-(1-Methyl-3-piperidyl)methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carboxamide oxalate;

N-[7-(2-Dimethylamino)ethoxy-2,3-dihydrobenzofuran-5-yl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-1,1'-biphenyl-4-carboxamide;

[3-(Dimethylaminoethyl)-3,6,7,8-tetrahydro-2H-furo[2,3-g]quinolin-5-yl]-[2'-methyl-4'-(5-methyl-[1,2,4]-oxadiazol-3-yl)-biphenyl-4-yl]-methanone oxalate;

5-[4-(2-Methyl-4-(5-methyl-1,2,4-oxadiazol-3-yl)phenyl)benzoylamino]-spiro[(2,3-dihydrobenzofuran)-3,4'-(1-methylhexahydroazepine)] oxalate;

2,3,5,6,7,8,-Hexahydro-1'-methyl-5-{2'-methyl-4'-[(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-yl]carbonyl}furo[2,3-g]quinoline-3-spiro-4'-piperidine oxalate;

6,7,8,9-Tetrahydro-1'-methyl-5-[[2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)][1,1'-biphenyl]-4-yl]carbonyl]-spiro[2H-furo[2,3-h][1]benzazepine-3(5H),4'-piperidine];

2,3,5,6,7,8-Hexahydro-5-[4'-(5-hydroxymethyl-1,2,4-oxadiazol-3-yl)-2'-methylbiphenyl-4-carbonyl]-1'-methylspiro[furo[2,3-g]quinoline-3,4'-piperidine] hydrochloride;

4'-(Dimethylamino)-5-[2'-methyl-4'-[(5-methyl-1,2,4-oxadiazol-3-yl)-biphenyl-4-yl]carbonyl]-2,3,6,7-tetrahydrospiro[furo[2,3-f]indole-3,1'-cyclohexane] hydrochloride;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-3-carboxamide;

N-[3-(3-Dimethylamino)propoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[4-(3-Dimethylamino)propoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[3-(Diethylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[3-[2-(Piperidin-1-yl)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-2-methylphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Dipropylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methylphenyl]-[1,1'-biphenyl]-4-carboxamide

N-[3-(2-Diisopropylamino)ethoxy-4-ethylphenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-4-(methoxycarbonylamino)phenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[1,1'-Biphenyl]-4-yl-3-(2-diisopropylamino)ethoxy-4-methoxybenzamide;

N-[3-(2-Diisopropylamino)ethoxy-4-(ethoxycarbonyl)phenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[4-(2-Diisopropylamino)ethoxy-3-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[2-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-diisopropylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[2-Fluoro-4-(2-diisopropylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[4-(2-Diisopropylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[2-(2-Diisopropylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4-bromobenzamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4-(isopropyl)benzamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4-(cyclohexyl)benzamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-3,4-dichlorobenzamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-3,5-dichlorobenzamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4-(nitro)benzamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-3-iodobenzamide;

N-[4-(2-Diisopropylamino)ethoxy-phenyl]-3-iodobenzamide;

N-[4-(2-Diisopropylamino)ethoxy-phenyl]-3-bromobenzamide;

N-[4-(2-Diisopropylamino)ethoxy-phenyl]-4-(cyclohexyl)benzamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxy-phenyl]-4-(cyclohexyl)benzamide;

N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-4'-ethyl-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-phenyl]-4'-ethyl-[1,1'-biphenyl]-4-carboxamide; and

N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-3'-methoxy-[1,1'-biphenyl]-4-carboxamide).

Among the most preferred compounds of the invention are the following compounds:

N-[3-(2-Diisopropylamino)ethoxy-phenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-(1,1'-biphenyl)-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-(1,1'-biphenyl)-4-carboxamide;

5-[4-(2-methyl-4-(5-methyl-1,2,4-oxadiazol-3-yl)phenyl)benzoylamino]spiro[(2,3-dihydrobenzofuran)-3,4'-(1-methylpiperidine)] oxalate;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-(1,1'-biphenyl)-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-2'-methyl-(1,1'-biphenyl)-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-phenyl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-[1,1'-biphenyl]-4-carboxamide;

N-[3-[2-(N-Cyclohexyl-N-isopropylamino)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-carboxamide;

N-[3-[2-(cis-2,6-Dimethylpiperidin-1-yl)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[3-[2-(N-Ethyl-N-isopropylamino)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[3-[2-(2,5-Dimethylpyrrolidin-1-yl)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[4-Methoxy-3-[2-(2,2,6,6-tetramethylpiperidin-1-yl)ethoxy]phenyl]-1,1'-biphenyl-4-carboxamide;

N-[3-[2-(N-t-Butyl-N-isopropyl)aminoethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[3-(3-Diisopropylamino)propyloxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[3-(3-Diisopropylamino)propyl-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-4-iodophenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-methanamine Dihydrochloride;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxy-phenyl]-4'-cyano-1,1'-biphenyl-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4'-cyano-2'-methyl-1,1'-biphenyl-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-2-naphthalenecarboxamide;

N-(1'-Methylspiro[benzofuran-3(2H),4'-piperidin]-5-yl-[1,1'-biphenyl]-4-carboxamide;

N-[3-[2-(N-Isopropyl-N-methylamino)ethoxy]-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4-(iodo)benzamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxy-phenyl]-4-(cyclohexyl)benzamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4'-ethyl-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-3'-methoxy-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-4'-acetyl-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-3'-methoxy-[1,1'-biphenyl]-4-carboxamide; and

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-3'-hydroxy-[1,1'-biphenyl]-4-carboxamide.

Compounds excluded from the scope of this invention are as follows:

N-[3-(2-Dimethylaminoethoxy)-4-methoxyphenyl]-4'-(5-dimethylamino-1,2,4-oxadiazol-3-yl)-2'-methylbiphenyl-4-carboxamide;

N-[3-(1-Methylazetidin-2-ylmethoxy)-4-methoxyphenyl]-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-2'-methylbiphenyl-4-carboxamide;

[7-(2-Dimethylaminoethoxy)-6-methoxy-3,4-dihydro-2H-quinolin-1-yl]-[2'-methyl-[1,2,4]-oxadiazol-3-yl]biphenyl-4-yl]-methanone oxalate;

2,3,6,7-Tetrahydro-1'-methyl-5-[2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carbonyl]-furo[2,3-f]indole-3-spiro-3'-piperidine oxalate;

5-[4'-(5-Methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carbonyl]-1'-methyl-2,3,6,7-tetrahydrofuro[2,3-f]indole-3-spiro-4'-piperidine;

1'-Ethyl-5-[2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)-biphenyl-4-carbonyl]-2,3,6,7-tetrahydrospirofuro[2,3-f]indole-3,4'-piperidine;

N-[2-(2-Dimethylamino)ethoxy-phenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]benzamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-2-carboxamide;

N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-4-(methoxy)benzamide;

N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-4-(nitro)benzamide;

N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-4-(nitro)benzamide;

N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-4-(nitro)benzamide;

N-[3-(Dimethylamino)ethoxy-4-methoxyphenyl]-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-4'-chloro-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-4'-hydroxy-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-4'-acetyl-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-4'-acetyl-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-3'-chloro-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-3'-nitro-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Dimethylamino)ethoxy-4-methoxyphenyl]-3'-nitro-[1,1'-biphenyl]-4-carboxamide;

N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-4-iodobenzamide;

N-[4-(2-Diethylamino)ethoxy-phenyl]-4-iodobenzamide;
 N-[2-(2-Diisopropylamino)ethoxy-phenyl]-4-iodobenzamide;
 N-[4-(2-Diethylamino)ethoxy-phenyl]-3-iodobenzamide;
 N-[2-(2-Diisopropylamino)ethoxy-phenyl]-3-iodobenzamide;
 N-[2-(2-Diethylamino)ethoxy-phenyl]-3-iodobenzamide;
 N-[3-(2-Diethylamino)ethoxy-phenyl]-4-bromobenzamide;
 N-[2-(2-Diisopropylamino)ethoxy-phenyl]-4-bromobenzamide;
 N-[2-(2-Diethylamino)ethoxy-phenyl]-4-bromobenzamide;
 N-[4-(2-Diethylamino)ethoxy-phenyl]-3-bromobenzamide;
 N-[2-(2-Diisopropylamino)ethoxy-phenyl]-3-bromobenzamide;
 N-[2-(2-Diethylamino)ethoxy-phenyl]-3-bromobenzamide;
 N-[3-(2-Diethylamino)ethoxy-4-methoxyphenyl]-4-(dimethylamino)benzamide;
 N-[4-(2-Diisopropylamino)ethoxy-phenyl]-4-(dimethylamino)benzamide;
 N-[4-(2-Diethylamino)ethoxy-phenyl]-4-(dimethylamino)benzamide;
 N-[2-(2-Diisopropylamino)ethoxy-phenyl]-4-(dimethylamino)benzamide;
 N-[2-(2-Diethylamino)ethoxy-phenyl]-4-(dimethylamino)benzamide;
 N-[4-(2-Diethylamino)ethoxy-phenyl]-4-(isopropyl)benzamide;
 N-[2-(2-Diisopropylamino)ethoxy-phenyl]-4-(isopropyl)benzamide;
 N-[2-(2-Diethylamino)ethoxy-phenyl]-4-(isopropyl)benzamide;
 N-[3-(2-Diethylamino)ethoxy-phenyl]-4-(cyclohexyl)benzamide;
 N-[2-(2-Diisopropylamino)ethoxy-phenyl]-4-(cyclohexyl)benzamide;
 N-[2-(2-Diethylamino)ethoxy-phenyl]-4-(cyclohexyl)benzamide;
 N-[2-(2-Diisopropylamino)ethoxy-phenyl]-4'-ethyl-[1,1'-biphenyl]-4-carboxamide; and
 N-[2-(2-Diethylamino)ethoxy-phenyl]-4'-ethyl-[1,1'-biphenyl]-4-carboxamide.

3. The method as claimed in claim 1, wherein the disease is selected from COPD, asthma and atopic disorders, rheumatoid arthritis, sarcoidosis and other fibrotic diseases, atherosclerosis, psoriasis, autoimmune diseases such as multiple sclerosis, inflammatory bowel disease, and HIV infection.

4. A compound selected from the group consisting of:

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-(1,1'-biphenyl)-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-2'-methyl-(1,1'-biphenyl)-4-carboxamide;

N-[3-(2-Diisopropylamino)ethoxy-4-methoxyphenyl]-2-naphthalenecarboxamide;

N-[2-(Dimethylaminomethyl)1,4-benzodioxan-7-yl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carboxamide; and

N-[2-(Dimethylaminomethyl)-1,4-benzodioxan-6-yl]-2'-methyl-4'-(5-methyl-1,2,4-oxadiazol-3-yl)biphenyl-4-carboxamide.

INTERNATIONAL SEARCH REPORT

International application No.
PCT/US98/13807

A. CLASSIFICATION OF SUBJECT MATTER

IPC(6) :A61K 31/41, 31/36, 31/135

US CL :514/364, 466, 651

According to International Patent Classification (IPC) or to both national classification and IPC

B. FIELDS SEARCHED

Minimum documentation searched (classification system followed by classification symbols)

U.S. : 514/364, 466, 651

Documentation searched other than minimum documentation to the extent that such documents are included in the fields searched

Electronic data base consulted during the international search (name of data base and, where practicable, search terms used)

C. DOCUMENTS CONSIDERED TO BE RELEVANT

Category*	Citation of document, with indication, where appropriate, of the relevant passages	Relevant to claim No.
A	US 3,931,195 A (DYKSTRA et al.) 06 January 1976.	1-4
A	US 4,000,143 A (DYKSTRA et al.) 28 December 1976	1-4
A,P	US 5,700,818 A (GASTER) 23 December 1997.	1-4

☐ Further documents are listed in the continuation of Box C.☐ See patent family annex.

* Special categories of cited documents:	*T* later document published after the international filing date or priority date and not in conflict with the application but cited to understand the principle or theory underlying the invention
A document defining the general state of the art which is not considered to be of particular relevance	*X* document of particular relevance; the claimed invention cannot be considered novel or cannot be considered to involve an inventive step when the document is taken alone
B earlier document published on or after the international filing date	*Y* document of particular relevance; the claimed invention cannot be considered to involve an inventive step when the document is combined with one or more other such documents, such combination being obvious to a person skilled in the art
L document which may throw doubts on priority claim(s) or which is cited to establish the publication date of another citation or other special reason (as specified)	*A* document member of the same patent family
O document referring to an oral disclosure, use, exhibition or other means	
P document published prior to the international filing date but later than the priority date claimed	

Date of the actual completion of the international search

21 AUGUST 1998

Date of mailing of the international search report

28 SEP 1998

Name and mailing address of the ISA/US
Commissioner of Patents and Trademarks
Box PCT
Washington, D.C. 20231

Facsimile No. (703) 305-3230

Authorized officer

RAYMOND J. HENLEY III

Telephone No. (703) 308-1235